

(12)

(21) 2 357 853

(22) 26.09.2001

(51) Int. Cl.⁷: **A61K 31/495, A61P 5/00,
A61K 31/41, A61K 31/435,
A61K 31/4427**

(30) 60/236,226 US 28.09.2000

(71) **PFIZER PRODUCTS INC.,**
Eastern Point Road, GROTON, XX (US).

(72) **FRYBURG, DAVID ALBERT (US).**


(74) **SMART & BIGGAR**

(54) UTILISATION DE SECRETAGOGUES DE L'HORMONE DE CROISSANCE CONJUGUEE A LA PRATIQUE DE L'EXERCICE

(54) USE OF GROWTH HORMONE SECRETAGOGUES IN CONJUNCTION WITH PHYSICAL EXERCISE

(57)

The present invention provides methods of using growth hormone secretagogues, prodrugs thereof and pharmaceutically acceptable salts of said secretagogues and said prodrugs, for increasing exercise-stimulated growth hormone secretion in a patient. The present invention also provides methods of using such compounds, in conjunction with physical exercise, to increase the improvement in muscle mass or strength in a patient. More specifically, the present invention provides such methods wherein the growth hormone secretagogues are compounds of Formula I: (see formula I) a prodrug thereof or a pharmaceutically acceptable salt of said secretagogue or said prodrug.





Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2357853 A1 2002/03/28

(21) 2 357 853

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(22) Date de dépôt/Filing Date: 2001/09/26

(41) Mise à la disp. pub./Open to Public Insp.: 2002/03/28

(30) Priorité/Priority: 2000/09/28 (60/236,226) US

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 31/495, A61K 31/4427,
A61K 31/435, A61K 31/41, A61P 5/00


(71) Demandeur/Applicant:
PFIZER PRODUCTS INC., US

(72) Inventeur/Inventor:
FRYBURG, DAVID ALBERT, US

(74) Agent: SMART & BIGGAR

(54) Titre : UTILISATION DE SECRETAGOGUES DE L'HORMONE DE CROISSANCE CONJUGUEE A LA PRATIQUE
DE L'EXERCICE

(54) Title: USE OF GROWTH HORMONE SECRETAGOGUES IN CONJUNCTION WITH PHYSICAL EXERCISE



(57) Abrégé/Abstract:

The present invention provides methods of using growth hormone secretagogues, prodrugs thereof and pharmaceutically acceptable salts of said secretagogues and said prodrugs, for increasing exercise-stimulated growth hormone secretion in a patient. The present invention also provides methods of using such compounds, in conjunction with physical exercise, to increase the improvement in muscle mass or strength in a patient. More specifically, the present invention provides such methods wherein the growth hormone secretagogues are compounds of Formula I: (see formula I) a prodrug thereof or a pharmaceutically acceptable salt of said secretagogue or said prodrug.

Canada

<http://opic.gc.ca> · Ottawa-Hull K1A 0C9 · <http://cipo.gc.ca>

OPIC · CIPO 191


OPIC



CIPO

ABSTRACT OF THE INVENTION

The present invention provides methods of using growth hormone secretagogues, prodrugs thereof and pharmaceutically acceptable salts of said secretagogues and said prodrugs, for increasing exercise-stimulated growth hormone secretion in a patient. The present invention also provides methods of using such compounds, in conjunction with physical exercise, to increase the improvement in muscle mass or strength in a patient. More specifically, the present invention provides such methods wherein the growth hormone secretagogues are compounds of Formula I:



10

a prodrug thereof or a pharmaceutically acceptable salt of said secretagogue or said prodrug.

72222-468

-1-

USE OF GROWTH HORMONE SECRETAGOGUES IN
CONJUNCTION WITH PHYSICAL EXERCISE
FIELD OF THE INVENTION

5 The present invention provides use of growth hormone
secretagogues, prodrugs thereof and pharmaceutically acceptable salts of the
secretagogues and the prodrugs, for increasing exercise-stimulated growth
hormone secretion in a patient. The present invention also provides use of
such compounds, together with physical exercise, to increase the
10 improvement in muscle mass or strength in a patient. More specifically, the
present invention provides such use wherein the growth hormone
secretagogues are certain compounds of Formula I below.

BACKGROUND OF THE INVENTION

Progressive decrease in muscle mass and strength (also known as
15 sarcopenia) are major determinants of physical frailty and loss of functional
independence in elderly or chronically ill individuals. Significant aging-associated loss
of muscle mass (also known as myopenia) and strength begins around the fifth
decade of life, but regular exercise, particularly high resistance exercise (strength
training), can largely prevent or attenuate this decline.

20 If a person does not exercise regularly, incremental gains are lost during
intervening periods of inactivity. Another problem is that the myopenia may be
caused in part by a shift in the set point in the muscle adaptive response, such that a
smaller mass gain is produced by the same amount of work in an elderly person.

Very few compounds are known in the art to be useful for the preservation
25 of muscle mass or strength produced by physical exercise. Anabolic steroids have
been used to increase muscle mass and enhance physical performance (*Anabolic-
Androgenic Steroids*, ed. Charles D. Kochakian, Vol. 43, Handbook of Experimental
Pharmacology (New York, Springer-Verlag, 1976) pp. 515-534). However, users of
anabolic steroids may suffer from side effects. An alternative way to enhance or
30 maintain the benefits of exercise would be highly desirable.

Growth hormone, which is secreted from the pituitary, stimulates growth of
all tissues of the body that are capable of growing. In addition, growth hormone is
known to have the following effects on the metabolic processes of the body: (1)
increased rate of protein synthesis in cells of the body; (2) decreased rate of
35 carbohydrate utilization in cells of the body; and (3) increased mobilization of free

fatty acids and use of fatty acids for energy. As is known to those skilled in the art, the known and potential uses of growth hormone are many and varied. See "Human Growth Hormone," Strobel and Thomas, Pharmacological Reviews, 46, pg. 1-34 (1994). Also, these varied uses of growth hormone are summarized in
5 International Patent Application, Publication Number WO 97/24369.

Various ways are known to release growth hormone (see Recent Progress in Hormone Research, vol. 52, pp. 215-245 (1997); and Front. Horm. Res., Basel, Karger, vol. 24, pp. 152-175 (1999)). For example, chemicals such as arginine, L-3,4-dihydroxyphenylalanine (L-DOPA), glucagon, vasopressin, and insulin induced
10 hypoglycemia, as well as activities such as sleep and exercise, indirectly cause growth hormone to be released from the pituitary by acting in some fashion on the hypothalamus perhaps either to decrease somatostatin secretion or to increase secretion of growth hormone releasing factor (GRF) or ghrelin (see Nature, vol. 402, pp. 656-660 (9 December 1999)), or all of these. Strobel and Thomas in
15 "Human Growth Hormone," which is referenced above, state that exercise is one of the most effective stimuli of growth hormone secretion.

In cases where increased levels of growth hormone were desired, the problem was generally solved by providing exogenous growth hormone or by administering GRF, insulin-like growth factor I (IGF-I) or a peptidyl compound which
20 stimulated growth hormone production and/or release. In any case, the peptidyl nature of the compound may limit the mode of administration, i.e., such compound may be administered, e.g., by injection or intranasally. Initially, the source of growth hormone was the extraction of the pituitary glands of cadavers. This resulted in a very expensive product and carried with it the risk that a disease associated with
25 the source of the pituitary gland could be transmitted to the recipient of the growth hormone. Recombinant growth hormone has become available which, while no longer carrying any such risk of disease transmission, is still a very expensive product which must be given by injection. In addition, administration of exogenous growth hormone may result in side-effects, including edema, and does not correlate
30 with the pulsatile release seen in the endogenous release of growth hormone.

Certain compounds have been developed which stimulate the release of endogenous growth hormone. Peptides which are known to stimulate the release of endogenous growth hormone include growth hormone releasing hormone and its analogs, the growth hormone releasing peptides, GHRP-6 and GHRP-1 (described

in U.S. Patent No. 4,411,890; International Patent Application, Publication No. WO 89/07110; and International Patent Application, Publication No. WO 89/07111), and GHRP-2 (described in International Patent Application, Publication No. WO 93/04081), as well as hexarelin (J. Endocrinol. Invest., 15 (Suppl. 4): 45 (1992)).

5 Other compounds possessing growth hormone secretagogue activity are disclosed in the following International Patent Applications (listed by Publication Nos.), issued U.S. Patents and published European Patent Applications: WO 98/46569, WO 98/51687, WO 98/58947, WO 98/58949, WO 98/58950, WO 99/08697, WO 99/09991, WO 95/13069, U.S. 5,492,916, U.S. 5,494,919, WO 95/14666, WO 10 94/19367, WO 94/13696, WO 94/11012, U.S. 5,726,319, WO 95/11029, WO 95/17422, WO 95/17423, WO 95/34311, WO 96/02530, WO 96/22996, WO 96/22997, WO 96/24580, WO 96/24587, U.S. 5,559,128, WO 96/32943, WO 96/33189, WO 96/15148, WO 96/38471, WO 96/35713, WO 97/00894, WO 97/07117, WO 97/06803, WO 97/11697, WO 97/15573, WO 97/22367, WO 15 97/23508, WO 97/22620, WO 97/22004, WO 97/21730, WO 97/24369, U.S. 5,663,171, WO 97/34604, WO 97/36873, WO 97/40071, WO 97/40023, WO 97/41878, WO97/41879, WO 97/46252, WO 97/44042, WO 97/38709, WO 98/03473, WO 97/43278, U.S. 5,721,251, U.S. 5,721,250, WO 98/10653, U.S. 5,919,777, U.S. 5,830,433 and EP 0995748.

20 In addition, the following growth hormone secretagogues are known in the art: MK-0677, L-162752 and L-163022 (Merck); NN703 and ipamorelin (Novo Nordisk); hexarelin (Pharmacia); GPA-748 (KP102, GHRP-2) (American Home Products); and LY444711 (Eli Lilly). The following agents that stimulate GH release via GHRH/GRF receptor (including GHRH/GRF derivatives, analogs and mimetics) 25 are known in the art: Geref (Ares/Serono); GHRH (1-44) (BioNebraska); Somatorelin (GRF 1-44) (Fujisawa/ICN); and ThGRF (Theratechnologies).

Endocrine Reviews 18(5): 621-645 (1997) provides an overview of peptidomimetic regulation of growth hormone secretion by growth hormone secretagogues. Horm. Res. 1999; 51(suppl 3):16-20 (1999), examines the clinical 30 and experimental effects of growth hormone secretagogues on various organ systems.

International Patent Applications, Publication Nos. WO 97/24369 and WO 98/58947, state that certain growth hormone secretagogues are useful for the treatment or prevention of osteoporosis, congestive heart failure, frailty associated

with aging, obesity, accelerating bone fracture repair, attenuating protein catabolic response after a major operation, reducing cachexia and protein loss due to chronic illness, accelerating wound healing, accelerating the recovery of burn patients, accelerating the recovery of patients having undergone major surgery, improving muscle strength, mobility, maintenance of skin thickness, metabolic homeostasis or renal homeostasis. Published European patent application 0995748 states that certain dipeptide growth hormone secretagogues are useful for the treatment or prevention of musculoskeletal frailty, including osteoporosis.

The administration of a growth hormone secretagogue is also known to enhance the quality of sleep, which is disclosed in International Patent Application, Publication No. WO 97/24369. Commonly assigned U.S. nonprovisional patent application 09/649622, filed 28 August 2000, provides pharmaceutical compositions comprising certain β_3 adrenergic agonists and growth hormone secretagogues or growth hormone, and their use for treating diabetes, obesity, hyperglycemia, frailty associated with obesity or frailty associated with aging, and for enhancing the quality of sleep in a mammal. International Patent Application, Publication No. WO 98/58949, discloses the treatment of insulin resistance with certain growth hormone secretagogues.

International Patent Application, Publication No. WO 00/12407, states that a growth hormone secretagogue is useful for enhancing the return of patients to independent living status following acute deconditioning such as that which may result from immobilization, surgery, or major injury such as hip fracture.

Journal of Orthopaedic Research 15:519:527 (1997) states that a growth hormone secretagogue, MK-677, elevated levels of serum insulin-like growth factor-I, which in turn increased the size and strength of the quadriceps muscle in canines during remobilization. J. Clin. Endocrinol. Metab. 83: 320-325, 1998, states that MK-677, an orally active growth hormone secretagogue, reverses diet-induced catabolism.

Drug Discovery Today, Vol. 4, No. 11, November 1999; and TEM Vol. 10, No. 1, 1999; discuss potential indications for growth hormone secretagogues, including their use in treating growth hormone disorders, such as growth hormone deficiency (GHD), age-related conditions, obesity and catabolic conditions, and their use in sleep enhancement.

72222-468

-5-

R.C. Cuneo et al., J. Appl. Physiol. 70(2): 695-700 (1991), studied the effects of growth hormone treatment on exercise performance in growth-hormone deficient adults.

R. Bross, M. Javanbakht and S. Bhasin, J. Clin.Endocrinol.Metab. 84:3420-3430 (1999), comment on anabolic intervention for the treatment of aging-associated sarcopenia.

A. Strawford et al., J.A.M.A, 281 (14):1282-1290 (1999), reported that recombinant growth hormone (rGH) and androgen replacement therapy in men with low or borderline serum testosterone concentrations are effective in restoring lean body mass in men with HIV infection.

Symposia Abstract No. 218, The Endocrine Society, 82nd Annual Meeting, Toronto, Ontario, Canada (June 21-21, 2000), investigated exercise and GHRH treatment of the elderly. Symposia Abstract No. 215, The Endocrine Society, 82nd Annual Meeting, Toronto, Ontario, Canada (June 21-21, 2000), investigated growth hormone and sex steroid treatment of the elderly.

H.C.M. Maas et al., Medicine & Science in Sports & Exercise, 32 (7): 1226-1232 (2000) investigated growth hormone responses during strenuous exercise and the role of GH-releasing hormone and GH-releasing peptide-2.

D.R. Taaffe and R. Marcus, Clinical Physiology (1997) 17, 311-324, investigated the effects of cessation and subsequent resumption of training on muscle strength in elderly men, just completing a 24-week trial of recombinant growth hormone (rhGH) and resistance exercise.

72222-468

-6-

SUMMARY OF THE INVENTION

The present invention provides drugs for increasing physical exercise-stimulated growth hormone secretion in a patient, which comprise a therapeutically effective amount of a growth hormone secretagogue. The present invention also provides such drugs that are so adapted to be taken before or at the time the patient performs physical exercise. Also, the present invention provides such drugs that are adapted to be taken between periods of physical exercise by the patient.

Also, the present invention provides drugs for increasing the improvement in physical performance produced by physical exercise in a patient, which comprise a therapeutically effective amount of a growth hormone secretagogue and are adapted to be taken in conjunction with physical exercise by the patient.

The present invention also provides drugs for increasing physical exercise capacity in a patient in need thereof, which comprise a therapeutically effective amount of a growth hormone secretagogue and are adapted to be taken in conjunction with physical exercise by the patient.

The present invention also provides drugs for maintaining or improving protein balance in a patient, which comprise a therapeutically effective amount of a growth hormone secretagogue and are adapted to be taken in conjunction with physical exercise by the patient.

72222-468

-7-

In addition, the present invention provides drugs for the preservation of the improvement in muscle mass or strength in a patient, which comprise a therapeutically effective amount of a growth hormone secretagogue, and are
5 adapted to be taken between periods of physical exercise by the patient.

The present invention also provides drugs for the preservation of the improvement in physical performance in a patient, which comprise a therapeutically effective amount
10 of a growth hormone secretagogue and are adapted to be taken between periods of physical exercise by the patient.

In addition, the present invention provides drugs for treating sarcopenia in a patient in need thereof, which comprise a therapeutically effective amount of a growth
15 hormone secretagogue and are adapted to be taken in conjunction with physical exercise by the patient.

More preferably, the present invention provides the above drugs wherein the physical exercise is physical rehabilitation or physical therapy. More preferably, the
20 present invention provides the above drugs wherein the physical exercise is part of a regular physical exercise program.

More preferably, the present invention provides the above drugs wherein the patient is a human. Even more
25 preferably, the present invention provides such drugs wherein the human is an elderly, chronically ill or obese individual. Also, the present invention provides such drugs wherein the human has age-related decline in physical performance or is growth hormone deficient.


72222-468

-7a-

More preferably, the present invention provides the above drugs wherein the growth hormone secretagogue is an orally active growth hormone secretagogue. Even more preferably, the present invention provides such drugs wherein the growth hormone secretagogue is orally administered.

More preferably, the growth hormone secretagogue is a compound of Formula I:


10



I

or a stereoisomeric mixture thereof, diastereomerically enriched, diastereomerically pure, enantiomerically enriched or enantiomerically pure isomer thereof, or a prodrug of such compound, mixture or isomer thereof, or a pharmaceutically acceptable salt of the compound, mixture, isomer or prodrug, or a tautomer thereof, wherein HET is a heterocyclic moiety selected from the group consisting of

-8-



d is 0, 1 or 2;

e is 1 or 2;

f is 0 or 1;

5 n and w are 0, 1 or 2, provided that n and w cannot both be 0 at the same time;

Y² is oxygen or sulfur;

A is a divalent radical, where the left hand side of the radical as shown below is connected to C'' and the right hand side of the radical as shown below is connected to C', selected from the group consisting of

- 10 -NR²-C(O)-NR²-, -NR²-S(O)₂-NR²-, -O-C(O)-NR²-, -NR²-C(O)-O-, -C(O)-NR²-C(O)-, -C(O)-NR²-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-NR²-C(O)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -S(O)₂-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-O-C(O)-, -C(R⁹R¹⁰)-O-C(R⁹R¹⁰)-, -NR²-C(O)-C(R⁹R¹⁰)-, -O-C(O)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(O)-NR²-, -C(R⁹R¹⁰)-C(O)-O-, -C(O)-NR²-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(O)-O-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -S(O)₂-NR²-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-NR²-C(O)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-O-C(O)-, -NR²-C(O)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -NR²-S(O)₂-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -O-C(O)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(O)-NR²-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(O)-, -C(R⁹R¹⁰)-NR²-C(O)-O-, -C(R⁹R¹⁰)-O-C(O)-NR²-, -C(R⁹R¹⁰)-NR²-C(O)-NR²-, -NR²-C(O)-O-C(R⁹R¹⁰)-, -NR²-C(O)-NR²-C(R⁹R¹⁰)-, -NR²-S(O)₂-NR²-C(R⁹R¹⁰)-, -O-C(O)-NR²-C(R⁹R¹⁰)-, -
- 15

- $C(O)-N=C(R^{11})-NR^2$ -, $-C(O)-NR^2-C(R^{11})=N$ -, $-C(R^9R^{10})-NR^{12}-C(R^9R^{10})$ -, $-NR^{12}-C(R^9R^{10})$ -, $-NR^{12}-C(R^9R^{10})-C(R^9R^{10})$ -, $-C(O)-O-C(R^9R^{10})-C(R^9R^{10})$ -, $-NR^2-C(R^{11})=N-C(O)$ -, $-C(R^9R^{10})-C(R^9R^{10})-N(R^{12})$ -, $-C(R^9R^{10})-NR^{12}$ -, $-N=C(R^{11})-NR^2-C(O)$ -, $-C(R^9R^{10})-C(R^9R^{10})-NR^2-S(O)_2$ -, $-C(R^9R^{10})-C(R^9R^{10})-S(O)_2-NR^2$ -, $-C(R^9R^{10})-C(R^9R^{10})-C(O)-O$ -, $-C(R^9R^{10})-S(O)_2-C(R^9R^{10})$ -, $-C(R^9R^{10})-C(R^9R^{10})-S(O)_2$ -, $-O-C(R^9R^{10})-C(R^9R^{10})$ -, $-C(R^9R^{10})-C(R^9R^{10})-O$ -, $-C(R^9R^{10})-C(O)-C(R^9R^{10})$ -, $-C(O)-C(R^9R^{10})-C(R^9R^{10})$ - and $-C(R^9R^{10})-NR^2-S(O)_2-NR^2$ -;
- Q is a covalent bond or CH_2 ;
- W is CH or N;
- 10 X is CR^9R^{10} , $C=CH_2$ or $C=O$;
- Y is CR^9R^{10} , O or NR^2 ;
- Z is $C=O$, $C=S$ or $S(O)_2$;
- G^1 is hydrogen, halo, hydroxy, nitro, amino, cyano, phenyl, carboxyl, $-CONH_2$, $-(C_1-C_4)$ alkyl optionally independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, $-(C_1-C_4)$ alkoxy optionally independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, $-(C_1-C_4)$ alkylthio, phenoxy, $-COO(C_1-C_4)$ alkyl, N,N-di- (C_1-C_4) alkylamino, $-(C_2-C_6)$ alkenyl optionally independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, $-(C_2-C_6)$ alkynyl optionally independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, $-(C_3-C_6)$ cycloalkyl optionally independently substituted with one or more (C_1-C_4) alkyl groups, one or more halogens or one or more hydroxy groups, $-(C_1-C_4)$ alkylamino carbonyl or di- (C_1-C_4) alkylamino carbonyl;
- G^2 and G^3 are each independently selected from the group consisting of hydrogen, halo, hydroxy, $-(C_1-C_4)$ alkyl optionally independently substituted with one to three halo groups and $-(C_1-C_4)$ alkoxy optionally independently substituted with one to three halo groups;
- R^1 is hydrogen, $-CN$, $-(CH_2)_qN(X^6)C(O)X^6$, $-(CH_2)_qN(X^6)C(O)(CH_2)_h-A^1$, $-(CH_2)_qN(X^6)S(O)_2(CH_2)_h-A^1$, $-(CH_2)_qN(X^6)S(O)_2X^6$, $-(CH_2)_qN(X^6)C(O)N(X^6)(CH_2)_h-A^1$, $-(CH_2)_qN(X^6)C(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)N(X^6)(CH_2)_h-A^1$, $-(CH_2)_qC(O)OX^6$, $-(CH_2)_qC(O)O(CH_2)_h-A^1$, $-(CH_2)_qOX^6$, $-(CH_2)_qOC(O)X^6$, $-(CH_2)_qOC(O)(CH_2)_h-A^1$, $-(CH_2)_qOC(O)N(X^6)(CH_2)_h-A^1$, $-(CH_2)_qOC(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)X^6$, $-(CH_2)_qC(O)(CH_2)_h-A^1$, $-(CH_2)_qN(X^6)C(O)OX^6$,
- 30

-10-

$-(CH_2)_qN(X^6)S(O)_2N(X^6)(X^6)$, $-(CH_2)_qS(O)_mX^6$, $-(CH_2)_qS(O)_m(CH_2)_tA^1$,
 $-(C_1-C_{10})alkyl$, $-(CH_2)_tA^1$, $-(CH_2)_q-(C_3-C_7)cycloalkyl$, $-(CH_2)_q-Y^1-(C_1-C_8)alkyl$,
 $-(CH_2)_q-Y^1-(CH_2)_tA^1$ or $-(CH_2)_q-Y^1-(CH_2)_t-(C_3-C_7)cycloalkyl$;

5 where the alkyl and cycloalkyl groups in the definition of R^1 are optionally substituted with $(C_1-C_4)alkyl$, hydroxy, $(C_1-C_4)alkoxy$, carboxyl, $-CONH_2$,
 $-S(O)_m(C_1-C_6)alkyl$, $-CO_2(C_1-C_4)alkyl$ ester, 1H-tetrazol-5-yl or 1, 2 or 3 fluoro groups;

Y^1 is O, $S(O)_m$, $-C(O)NX^6$ -, $-CH=CH-$, $-C\equiv C-$, $-N(X^6)C(O)-$, $-C(O)NX^6$ -,
 $-C(O)O-$, $-OC(O)N(X^6)-$ or $-OC(O)-$;

10 q is 0, 1, 2, 3 or 4;

t is 0, 1, 2 or 3;

said $(CH_2)_q$ group and $(CH_2)_t$ group in the definition of R^1 are optionally independently substituted with hydroxy, $(C_1-C_4)alkoxy$, carboxyl, $-CONH_2$,
 $-S(O)_m(C_1-C_6)alkyl$, $-CO_2(C_1-C_4)alkyl$ ester, 1H-tetrazol-5-yl, 1, 2 or 3 fluoro groups or 1 or 2 $(C_1-C_4)alkyl$ groups;

15 R^{1A} is selected from the group consisting of hydrogen, F, Cl, Br, I, $(C_1-C_6)alkyl$, phenyl $(C_1-C_3)alkyl$, pyridyl $(C_1-C_3)alkyl$, thiazolyl $(C_1-C_3)alkyl$ and thienyl $(C_1-C_3)alkyl$, provided that R^{1A} is not F, Cl, Br or I when a heteroatom is vicinal to C^* ;

R^2 is hydrogen, $(C_1-C_6)alkyl$, $-(C_0-C_3)alkyl-(C_3-C_8)cycloalkyl$, $-(C_1-C_4)alkyl-A^1$ or A^1 ;

20 where the alkyl groups and the cycloalkyl groups in the definition of R^2 are optionally substituted with hydroxy, $-C(O)OX^6$, $-C(O)N(X^6)(X^6)$, $-N(X^6)(X^6)$, $-S(O)_m(C_1-C_6)alkyl$, $-C(O)A^1$, $-C(O)(X^6)$, CF_3 , CN or 1, 2 or 3 independently selected halo groups;

25 R^3 is selected from the group consisting of A^1 , $(C_1-C_{10})alkyl$, $-(C_1-C_6)alkyl-A^1$, $-(C_1-C_6)alkyl-(C_3-C_7)cycloalkyl$, $-(C_1-C_5)alkyl-X^1-(C_1-C_5)alkyl$, $-(C_1-C_5)alkyl-X^1-(C_0-C_5)alkyl-A^1$ and $-(C_1-C_5)alkyl-X^1-(C_1-C_5)alkyl-(C_3-C_7)cycloalkyl$;

where the alkyl groups in the definition of R^3 are optionally substituted with $-S(O)_m(C_1-C_6)alkyl$, $-C(O)OX^3$, 1, 2, 3, 4 or 5 independently selected halo groups or 1, 2 or 3 independently selected $-OX^3$ groups;


30 X^1 is O, $S(O)_m$, $-N(X^2)C(O)-$, $-C(O)N(X^2)-$, $-OC(O)-$, $-C(O)O-$, $-CX^2=CX^2-$, $-N(X^2)C(O)O-$, $-OC(O)N(X^2)-$ or $-C\equiv C-$;

R^4 is hydrogen, $(C_1-C_6)alkyl$ or $(C_3-C_7)cycloalkyl$, or R^4 is taken together with R^3 and the carbon atom to which they are attached and form $(C_5-C_7)cycloalkyl$, $(C_5-C_7)cycloalkenyl$, a partially saturated or fully saturated 4- to 8-membered ring

-11-

having 1 to 4 heteroatoms independently selected from the group consisting of oxygen, sulfur and nitrogen, or is a bicyclic ring system consisting of a partially saturated or fully saturated 5- or 6-membered ring, fused to a partially saturated, fully unsaturated or fully saturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

X^4 is hydrogen or (C_1-C_6) alkyl or X^4 is taken together with R^4 and the nitrogen atom to which X^4 is attached and the carbon atom to which R^4 is attached and form a five to seven membered ring;



where a and b are each independently 0, 1, 2 or 3;

X^5 and X^{5a} are each independently selected from the group consisting of hydrogen, CF_3 , A^1 and optionally substituted (C_1-C_6) alkyl;

15 the optionally substituted (C_1-C_6) alkyl in the definition of X^5 and X^{5a} is optionally substituted with a substituent selected from the group consisting of A^1 , OX^2 , $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^2$, (C_3-C_7) cycloalkyl, $-N(X^2)(X^2)$ and $-C(O)N(X^2)(X^2)$;

20 or the carbon bearing X^5 or X^{5a} forms one or two alkylene bridges with the nitrogen atom bearing R^7 and R^8 wherein each alkylene bridge contains 1 to 5 carbon atoms, provided that when one alkylene bridge is formed then only one of X^5 or X^{5a} is on the carbon atom and only one of R^7 or R^8 is on the nitrogen atom and further provided that when two alkylene bridges are formed then X^5 and X^{5a} cannot be on the carbon atom and R^7 and R^8 cannot be on the nitrogen atom;

25 or X^5 is taken together with X^{5a} and the carbon atom to which they are attached and form a partially saturated or fully saturated 3- to 7-membered ring, or a partially saturated or fully saturated 4- to 8-membered ring having 1 to 4 heteroatoms independently selected from the group consisting of oxygen, sulfur and nitrogen;

30 or X^5 is taken together with X^{5a} and the carbon atom to which they are attached and form a bicyclic ring system consisting of a partially saturated or fully saturated 5- or 6-membered ring, optionally having 1 or 2


-12-

heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated, fully saturated or fully unsaturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

Z^1 is a bond, O or $N-X^2$, provided that when a and b are both 0 then Z^1 is not $N-X^2$ or O;

or R^b is $-(CR^aR^b)_a-E-(CR^aR^b)_b-$, where the $-(CR^aR^b)_a-$ group is attached to the carbonyl carbon of the amide group of the compound of formula I and the $-(CR^aR^b)_b$ group is attached to the terminal nitrogen atom of the compound of Formula I;

E is -O-, -S-, -CH=CH- or an aromatic moiety selected from



said aromatic moiety in the definition of E optionally substituted with up to three halo, hydroxy, $-N(R^c)(R^c)$, (C_1-C_6) alkyl or (C_1-C_6) alkoxy;

R^a and R^b are, for each occurrence, independently hydrogen, (C_1-C_6) alkyl, trifluoromethyl, phenyl or monosubstituted (C_1-C_6) alkyl where the substituents are imidazolyl, naphthyl, phenyl, indolyl, p-hydroxyphenyl, $-OR^c$, $S(O)_mR^c$, $C(O)OR^c$, (C_3-C_7) cycloalkyl, $-N(R^c)(R^c)$, $-C(O)N(R^c)(R^c)$, or R^a or R^b may independently be joined to one or both of R^7 or E (where E is other than O, S or -CH=CH-) to form an alkylene bridge between the terminal nitrogen and the alkyl portion of the R^a or R^b and the R^7 or E group, wherein the bridge contains 1 to 8 carbon atoms; or R^a and R^b may be joined to one another to form a (C_3-C_7) cycloalkyl;

R^c , for each occurrence, is independently hydrogen or (C_1-C_6) alkyl; a and b are independently 0, 1, 2 or 3, with the proviso that if E is -O- or -S-, b is other than 0 or 1 and with the further proviso that if E is -CH=CH-, b is other than 0;

R^7 and R^8 are each independently hydrogen or optionally substituted (C_1-C_6) alkyl;
 where the optionally substituted (C_1-C_6) alkyl in the definition of R^7 and R^8 is
 optionally independently substituted with A^1 , $-C(O)O-(C_1-C_6)$ alkyl,
 $-S(O)_m(C_1-C_6)$ alkyl, 1 to 5 halo groups, 1 to 3 hydroxy groups, 1 to 3
 5 $-O-C(O)(C_1-C_{10})$ alkyl groups or 1 to 3 (C_1-C_6) alkoxy groups; or

R^7 and R^8 can be taken together to form $-(CH_2)_r-L-(CH_2)_r-$;

where L is $C(X^2)(X^2)$, $S(O)_m$ or $N(X^2)$;

R^9 and R^{10} are each independently selected from the group consisting of hydrogen,
 fluoro, hydroxy and (C_1-C_5) alkyl optionally independently substituted with 1-5 halo
 10 groups;

R^{11} is selected from the group consisting of (C_1-C_5) alkyl and phenyl optionally
 substituted with 1-3 substituents each independently selected from the group
 consisting of (C_1-C_5) alkyl, halo and (C_1-C_5) alkoxy;

R^{12} is selected from the group consisting of (C_1-C_5) alkylsulfonyl, (C_1-C_5) alkanoyl
 15 and (C_1-C_5) alkyl where the alkyl portion is optionally independently substituted by 1-
 5 halo groups;

A^1 for each occurrence is independently selected from the group consisting of $(C_5-$
 $C_7)$ cycloalkenyl, phenyl, a partially saturated, fully saturated or fully unsaturated 4-
 to 8-membered ring optionally having 1 to 4 heteroatoms independently selected
 20 from the group consisting of oxygen, sulfur and nitrogen and a bicyclic ring system
 consisting of a partially saturated, fully unsaturated or fully saturated 5- or 6-
 membered ring, optionally having 1 to 4 heteroatoms independently selected from
 the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated,
 fully saturated or fully unsaturated 5- or 6-membered ring, optionally having 1 to 4
 25 heteroatoms independently selected from the group consisting of nitrogen, sulfur
 and oxygen;

A^1 for each occurrence is independently optionally substituted, on one or
 optionally both rings if A^1 is a bicyclic ring system, with up to three
 substituents, each substituent independently selected from the group
 30 consisting of F, Cl, Br, I, OCF_3 , OCF_2H , CF_3 , CH_3 , OCH_3 , $-OX^6$,
 $-C(O)N(X^6)(X^6)$, $-C(O)OX^6$, oxo, (C_1-C_6) alkyl, nitro, cyano, benzyl, $-$
 $S(O)_m(C_1-C_6)$ alkyl, 1H-tetrazol-5-yl, phenyl, phenoxy, phenylalkyloxy,
 halophenyl, methylenedioxy, $-N(X^6)(X^6)$, $-N(X^6)C(O)(X^6)$, $-S(O)_2N(X^6)(X^6)$,
 $-N(X^6)S(O)_2$ -phenyl, $-N(X^6)S(O)_2X^6$, $-CONX^{11}X^{12}$, $-S(O)_2NX^{11}X^{12}$,

-14-

$-NX^6S(O)_2X^{12}$, $-NX^6CONX^{11}X^{12}$, $-NX^6S(O)_2NX^{11}X^{12}$, $-NX^6C(O)X^{12}$, imidazolyl, thiazolyl and tetrazolyl, provided that if A^1 is optionally substituted with methylenedioxy then it can only be substituted with one methylenedioxy;

where X^{11} is hydrogen or optionally substituted (C_1-C_6) alkyl;

5 the optionally substituted (C_1-C_6) alkyl defined for X^{11} is optionally independently substituted with phenyl, phenoxy, (C_1-C_6) alkoxycarbonyl, $-S(O)_m(C_1-C_6)$ alkyl, 1 to 5 halo groups, 1 to 3 hydroxy groups, 1 to 3 (C_1-C_{10}) alkanoyloxy groups or 1 to 3 (C_1-C_6) alkoxy groups;

10 X^{12} is hydrogen, (C_1-C_6) alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl, provided that when X^{12} is not hydrogen, the X^{12} group is optionally substituted with one to three substituents independently selected from the group consisting of Cl, F, CH_3 , OCH_3 , OCF_3 and CF_3 ;

15 or X^{11} and X^{12} are taken together to form $-(CH_2)_r-L^1-(CH_2)_r-$;

L^1 is $C(X^2)(X^2)$, O, $S(O)_m$ or $N(X^2)$;

r for each occurrence is independently 1, 2 or 3;

X^2 for each occurrence is independently hydrogen, optionally substituted (C_1-C_6) alkyl or optionally substituted (C_3-C_7) cycloalkyl, where the optionally substituted (C_1-C_6) alkyl and optionally substituted (C_3-C_7) cycloalkyl in the definition of X^2 are
20 optionally independently substituted with $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^3$, 1 to 5 halo groups or 1-3 OX^3 groups;

X^3 for each occurrence is independently hydrogen or (C_1-C_6) alkyl;

X^6 for each occurrence is independently hydrogen, optionally substituted (C_1-C_6) alkyl, (C_2-C_6) halogenated alkyl, optionally substituted (C_3-C_7) cycloalkyl, (C_3-C_7) -
25 halogenated cycloalkyl, where optionally substituted (C_1-C_6) alkyl and optionally substituted (C_3-C_7) cycloalkyl in the definition of X^6 is optionally independently mono- or di-substituted with (C_1-C_4) alkyl, hydroxy, (C_1-C_4) alkoxy, carboxyl, $CONH_2$, $-S(O)_m(C_1-C_6)$ alkyl, carboxylate (C_1-C_4) alkyl ester or 1H-tetrazol-5-yl; or

30 when there are two X^6 groups on one atom and both X^6 are independently (C_1-C_6) alkyl, the two (C_1-C_6) alkyl groups may be optionally joined and, together with the atom to which the two X^6 groups are attached, form a 4- to 9- membered ring optionally having oxygen, sulfur or NX^7 as a ring member;

X^7 is hydrogen or (C_1-C_6) alkyl optionally substituted with hydroxy;

-15-


m for each occurrence is independently 0, 1 or 2;

with the provisos that:

1) X^6 and X^{12} cannot be hydrogen when attached to $C(O)$ or $S(O)_2$ in the form $C(O)X^6$, $C(O)X^{12}$, $S(O)_2X^6$ or $S(O)_2X^{12}$; and

5 2) when R^6 is a bond then L is $N(X^2)$ and each r in the definition $-(CH_2)_r-L-(CH_2)_r$ is independently 2 or 3.

More preferably, the present invention provides such methods wherein the growth hormone secretagogue is a compound having the structural formula below, which is designated herein as Formula I-A



a racemic-diastereomeric mixture or optical isomer of said compound or a pharmaceutically-acceptable salt or a prodrug thereof, or a tautomer thereof, wherein

15 f is 0;

n is 0 and w is 2, or n is 1 and w is 1, or n is 2 and w is 0;

Y is oxygen or sulfur;

R^1 is hydrogen, $-CN$, $-(CH_2)_qN(X^6)C(O)X^6$, $-(CH_2)_qN(X^6)C(O)(CH_2)_t-A^1$,

$-(CH_2)_qN(X^6)SO_2(CH_2)_t-A^1$, $-(CH_2)_qN(X^6)SO_2X^6$, $-(CH_2)_qN(X^6)C(O)N(X^6)(CH_2)_t-A^1$,

20 $-(CH_2)_qN(X^6)C(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)N(X^6)(CH_2)_t-A^1$,

$-(CH_2)_qC(O)OX^6$, $-(CH_2)_qC(O)O(CH_2)_t-A^1$, $-(CH_2)_qOX^6$, $-(CH_2)_qOC(O)X^6$,

$-(CH_2)_qOC(O)(CH_2)_t-A^1$, $-(CH_2)_qOC(O)N(X^6)(CH_2)_t-A^1$, $-(CH_2)_qOC(O)N(X^6)(X^6)$,

$-(CH_2)_qC(O)X^6$, $-(CH_2)_qC(O)(CH_2)_t-A^1$, $-(CH_2)_qN(X^6)C(O)OX^6$,

$-(CH_2)_qN(X^6)SO_2N(X^6)(X^6)$, $-(CH_2)_qS(O)_mX^6$, $-(CH_2)_qS(O)_m(CH_2)_t-A^1$,

25 $-(C_1-C_{10})$ alkyl, $-(CH_2)_t-A^1$, $-(CH_2)_q-(C_3-C_7)$ cycloalkyl, $-(CH_2)_q-Y^1-(C_1-C_6)$ alkyl,

$-(CH_2)_q-Y^1-(CH_2)_t-A^1$ or $-(CH_2)_q-Y^1-(CH_2)_t-(C_3-C_7)$ cycloalkyl;

where the alkyl and cycloalkyl groups in the definition of R^1 are optionally substituted with (C_1-C_4) alkyl, hydroxyl, (C_1-C_4) alkoxy, carboxyl, $-CONH_2$,

$-S(O)_m(C_1-C_6)$ alkyl, $-CO_2(C_1-C_4)$ alkyl ester, 1H-tetrazol-5-yl or 1, 2 or 3

30

fluoro;

-16-

Y^1 is O, $S(O)_m$, $-C(O)NX^6$ -, $-CH=CH$ -, $-C\equiv C$ -, $-N(X^6)C(O)$ -, $-C(O)O$ -,
 $-OC(O)N(X^6)$ - or $-OC(O)$ -;

q is 0, 1, 2, 3 or 4;

t is 0, 1, 2 or 3;

- 5 said $(CH_2)_q$ group and $(CH_2)_t$ group may each be optionally substituted with hydroxyl, (C_1-C_4) alkoxy, carboxyl, $-CONH_2$, $-S(O)_m(C_1-C_6)$ alkyl, $-CO_2(C_1-C_4)$ alkyl ester, 1H-tetrazol-5-yl, 1, 2 or 3 fluoro, or 1 or 2 (C_1-C_4) alkyl;

R^2 is hydrogen, (C_1-C_8) alkyl, $-(C_0-C_3)$ alkyl- (C_3-C_8) cycloalkyl, $-(C_1-C_4)$ alkyl- A^1 or A^1 ;

- 10 where the alkyl groups and the cycloalkyl groups in the definition of R^2 are optionally substituted with hydroxyl, $-C(O)OX^6$, $-C(O)N(X^6)(X^6)$, $-N(X^6)(X^6)$, $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)A^1$, $-C(O)(X^6)$, CF_3 , CN or 1, 2 or 3 halogen;

R^3 is A^1 , (C_1-C_{10}) alkyl, $-(C_1-C_6)$ alkyl- A^1 , $-(C_1-C_6)$ alkyl- (C_3-C_7) cycloalkyl,

- 15 $-(C_1-C_5)$ alkyl- X^1 -(C_1-C_5)alkyl, $-(C_1-C_5)$ alkyl- X^1 -(C_0-C_5)alkyl- A^1 or $-(C_1-C_5)$ alkyl- X^1 -(C_1-C_5)alkyl- (C_3-C_7) cycloalkyl;

where the alkyl groups in the definition of R^3 are optionally substituted with,


$-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^3$, 1, 2, 3, 4 or 5 halogens, or 1, 2 or 3 OX^3 ;

X^1 is O, $S(O)_m$, $-N(X^2)C(O)$ -, $-C(O)N(X^2)$ -, $-OC(O)$ -, $-C(O)O$ -, $-CX^2=CX^2$ -,

- 20 $-N(X^2)C(O)O$ -, $-OC(O)N(X^2)$ - or $-C\equiv C$;

R^4 is hydrogen, (C_1-C_6) alkyl or (C_3-C_7) cycloalkyl;

X^4 is hydrogen or (C_1-C_6) alkyl or X^4 is taken together with R^4 and the nitrogen atom to which X^4 is attached and the carbon atom to which R^4 is attached and form a five to seven membered ring;



where a and b are independently 0, 1, 2 or 3;

X^5 and X^{5a} are each independently selected from the group consisting of hydrogen, trifluoromethyl, A^1 and optionally substituted (C_1-C_6) alkyl;

the optionally substituted (C_1-C_6) alkyl in the definition of X^5 and X^{5a} is

- 30 optionally substituted with a substituent selected from the group consisting of A^1 , OX^2 , $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^2$, (C_3-C_7) cycloalkyl, $-N(X^2)(X^2)$ and $-C(O)N(X^2)(X^2)$;

R^7 and R^8 are independently hydrogen or optionally substituted (C_1-C_6)alkyl;
 where the optionally substituted (C_1-C_6)alkyl in the definition of R^7 and R^8 is
 optionally independently substituted with A^1 , $-C(O)O-(C_1-C_6)$ alkyl,
 $-S(O)_m(C_1-C_6)$ alkyl, 1 to 5 halogens, 1 to 3 hydroxy, 1 to 3 $-O-C(O)(C_1-$
 5 $C_{10})$ alkyl or 1 to 3 (C_1-C_6)alkoxy; or

R^7 and R^8 can be taken together to form $-(CH_2)_i-L-(CH_2)_i-$;
 where L is $C(X^2)(X^2)$, $S(O)_m$ or $N(X^2)$;

A^1 in the definition of R^1 is a partially saturated, fully saturated or fully unsaturated
 4- to 8-membered ring optionally having 1 to 4 heteroatoms independently selected
 10 from the group consisting of oxygen, sulfur and nitrogen, a bicyclic ring system
 consisting of a partially saturated, fully unsaturated or fully saturated 5- or 6-
 membered ring, having 1 to 4 heteroatoms independently selected from the group
 consisting of nitrogen, sulfur and oxygen, fused to a partially saturated, fully
 saturated or fully unsaturated 5- or 6-membered ring, optionally having 1 to 4
 15 heteroatoms independently selected from the group consisting of nitrogen, sulfur
 and oxygen;

A^1 in the definition of R^2 , R^3 , R^6 , R^7 and R^8 is independently (C_5-C_7)cycloalkenyl,
 phenyl or a partially saturated, fully saturated or fully unsaturated 4- to 8-
 membered ring optionally having 1 to 4 heteroatoms independently selected from
 20 the group consisting of oxygen, sulfur and nitrogen, a bicyclic ring system
 consisting of a partially saturated, fully unsaturated or fully saturated 5- or 6-
 membered ring, optionally having 1 to 4 heteroatoms independently selected from
 the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated,
 fully saturated or fully unsaturated 5- or 6- membered ring, optionally having 1 to 4
 25 heteroatoms independently selected from the group consisting of nitrogen, sulfur
 and oxygen;

A^1 for each occurrence is independently optionally substituted, in one or
 optionally both rings if A^1 is a bicyclic ring system, with up to three
 substituents, each substituent independently selected from the group
 30 consisting of F, Cl, Br, I, OCF_3 , OCF_2H , CF_3 , CH_3 , OCH_3 , $-OX^6$,
 $-C(O)N(X^6)(X^6)$, $-C(O)OX^6$, oxo, (C_1-C_6)alkyl, nitro, cyano, benzyl,
 $-S(O)_m(C_1-C_6)$ alkyl, 1H-tetrazol-5-yl, phenyl, phenoxy, phenylalkyloxy,
 halophenyl, methylenedioxy, $-N(X^6)(X^6)$, $-N(X^6)C(O)(X^6)$, $-SO_2N(X^6)(X^6)$,
 $-N(X^6)SO_2$ -phenyl, $-N(X^6)SO_2X^6$, $-CONX^{11}X^{12}$, $-SO_2NX^{11}X^{12}$, $-NX^6SO_2X^{12}$,

$-NX^6CONX^{11}X^{12}$, $-NX^6SO_2NX^{11}X^{12}$, $-NX^6C(O)X^{12}$, imidazolyl, thiazolyl or tetrazolyl, provided that if A^1 is optionally substituted with methylenedioxy then it can only be substituted with one methylenedioxy;

where X^{11} is hydrogen or optionally substituted (C_1-C_6) alkyl;

5 the optionally substituted (C_1-C_6) alkyl defined for X^{11} is optionally independently substituted with phenyl, phenoxy, (C_1-C_6) alkoxycarbonyl, $-S(O)_m(C_1-C_6)$ alkyl 1 to 5 halogens, 1 to 3 hydroxy, 1 to 3 (C_1-C_{10}) alkanoyloxy or 1 to 3 (C_1-C_6) alkoxy;

10 X^{12} is hydrogen, (C_1-C_6) alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl, provided that when X^{12} is not hydrogen, X^{12} is optionally substituted with one to three substituents independently selected from the group consisting of Cl, F, CH_3 , OCH_3 , OCF_3 and CF_3 ;

or X^{11} and X^{12} are taken together to form $-(CH_2)_rL^1-(CH_2)_r$;

15 where L^1 is $C(X^2)(X^2)$, O, $S(O)_m$ or $N(X^2)$;

r for each occurrence is independently 1, 2 or 3;

X^2 for each occurrence is independently hydrogen, optionally substituted (C_1-C_6) alkyl, or optionally substituted (C_3-C_7) cycloalkyl, where the optionally substituted (C_1-C_6) alkyl and optionally substituted (C_3-C_7) cycloalkyl in the definition of X^2 are
20 optionally independently substituted with $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^3$, 1 to 5 halogens or 1-3 OX^3 ;

X^3 for each occurrence is independently hydrogen or (C_1-C_6) alkyl;

X^6 is independently hydrogen, optionally substituted (C_1-C_6) alkyl, (C_2-C_6) halogenated alkyl, optionally substituted (C_3-C_7) cycloalkyl, (C_3-C_7) -
25 halogenatedcycloalkyl, where optionally substituted (C_1-C_6) alkyl and optionally substituted (C_3-C_7) cycloalkyl in the definition of X^6 is optionally independently substituted by 1 or 2 (C_1-C_4) alkyl, hydroxyl, (C_1-C_4) alkoxy, carboxyl, $CONH_2$, $-S(O)_m(C_1-C_6)$ alkyl, carboxylate (C_1-C_4) alkyl ester, or 1H-tetrazol-5-yl; or

when there are two X^6 groups on one atom and both X^6 are independently (C_1-C_6) alkyl, the two (C_1-C_6) alkyl groups may be optionally joined and, together with the
30 atom to which the two X^6 groups are attached, form a 4- to 9- membered ring optionally having oxygen, sulfur or NX^7 ;

X^7 is hydrogen or (C_1-C_6) alkyl optionally substituted with hydroxyl; and

m for each occurrence is independently 0, 1 or 2;

72222-468

-19-

with the proviso that:

X^6 and X^{12} cannot be hydrogen when it is attached to C(O) or SO_2 in the form $C(O)X^6$, $C(O)X^{12}$, SO_2X^6 or SO_2X^{12} ; and

when R^6 is a bond then L is $N(X^2)$ and each r in the definition $-(CH_2)_r-L-(CH_2)_r-$ is

5 independently 2 or 3.

More preferably, the present invention provides such drugs wherein the growth hormone secretagogue is 2-amino-N-(2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl)-isobutyramide, a prodrug thereof or a pharmaceutically acceptable salt of the growth hormone secretagogue or the prodrug. Even more preferably, the present invention provides such drugs wherein the growth hormone secretagogue is 2-amino-N-[2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl]-isobutyramide, L-tartrate.

Also, more preferably, the present invention provides such drugs wherein the growth hormone secretagogue is 2-amino-N-(1-(R)-(2,4-difluoro-benzyloxymethyl)-2-oxo-2-(3-oxo-3a-(R)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-ethyl)-2-methyl-propionamide, a prodrug thereof or a pharmaceutically acceptable salt of the growth hormone secretagogue or the prodrug. Even more preferably, the present invention provides such drugs wherein the growth hormone secretagogue is the (L)-(+)-tartaric acid salt of 2-amino-N-(1-(R)-(2,4-difluoro-benzyloxymethyl)-2-oxo-2-(3-oxo-3a-(R)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-ethyl)-2-methyl-propionamide.

Also, more preferably, the present invention provides such drugs wherein the growth hormone secretagogue is 2-amino-N-(1-(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-hexahydro-imidazo[1,5-a]pyrazin-7-yl)-2-oxo-ethyl)-2-methyl-propionamide, a prodrug thereof or a pharmaceutically acceptable salt of the growth hormone secretagogue or the prodrug. Even more preferably, the present invention provides such drugs wherein the growth hormone secretagogue is the (L)-(+)-tartaric acid salt of 2-amino-N-(1-(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-hexahydro-imidazo[1,5-a]pyrazin-7-yl)-2-oxo-ethyl)-2-methyl-propionamide.

72222-468

-20-

The present invention further provides drugs, as described above, which further comprise recombinant growth hormone or a growth hormone secretagogue selected from the group consisting of GHRP-6, GHRP-1, GHRP-2, hexarelin, growth hormone releasing factor, an analog of growth hormone releasing factor, IGF-I and IGF-II.

The present invention further provides drugs, as described above, which further comprise an anabolic steroid.

The present invention further provides drugs, as described above, which further comprise a naturaceutic.

The present invention further provides commercial packages or kits that comprise the above-mentioned drugs and written matters which are associated therewith and describe indications, directions or instructions of use of the drugs.

DETAILED DESCRIPTION OF THE INVENTION

The term "drug" in this specification means a pharmaceutical composition or formulation containing one or more pharmaceutically acceptable carriers in addition to the growth hormone secretagogue.

The present invention is directed to the use of a compound, which has the ability to stimulate or amplify the release of natural or endogenous growth hormone, to increase growth hormone secretion in response to exercise. Also, the present invention provides a drug for increasing the improvement in muscle mass or strength produced by physical exercise, comprising a growth hormone secretagogue,

72222-468

-20a-

preferably a compound of Formula I below. In addition, the present invention provides a drug for preserving or maintaining the improvement in muscle mass or strength, with or between bouts of exercise, comprising a growth hormone
5 secretagogue, preferably a compound of Formula I below.

Further, according to the present invention, a growth hormone secretagogue administered together with physical exercise may augment growth hormone release and its associated benefits. It may have beneficial action at the
10 level of muscle tissue, i.e., it may, for example, have a beneficial effect on amino acid transport or function of growth factors or their receptors in muscle tissue. It may promote muscle hypertrophy and preserve the gain in muscle mass and strength between periods of physical exercise. It
15 may maintain the homeostatic set point for muscle as a function of an individual's activity/exercise level. It may maintain or improve protein balance between periods of physical exercise. Breakdown of protein (protein catabolism), which occurs as a result of physical exercise,
20 may be measured by urinary nitrogen excretion.

The benefits of physical exercise are many and varied and are applicable to both young and old, and men and women. It is especially important for older

people to exercise because it helps to maintain physical and mental functioning in older individuals.

There are many forms of physical exercise. Physical exercise includes, for example, physical rehabilitation and physical therapy. It includes activities such as walking, running, jogging, aerobics, rowing, weight-lifting, swimming, and many others known in the art. Physical exercise may be performed occasionally or on a regular basis, depending on the physical requirements and/or condition of the person who is exercising.

In a preferred embodiment of the present invention, therapy with the growth hormone secretagogue is supplemented or employed together with physical exercise, such as physical rehabilitation. When employed in conjunction with physical exercise, the desired secretory effect on growth hormone may be achieved by using a lower dose of the growth hormone secretagogue. Thus, certain side effects of growth hormone secretagogue treatment may be lessened.

According to the present invention, the growth hormone secretagogue is preferably administered in conjunction with physical exercise by the patient. Thus, the growth hormone secretagogue may be administered before the patient performs physical exercise, during the time when the patient is performing physical exercise or after the patient has performed physical exercise, or any combination of any of these time periods. It may also be administered between periods of physical exercise by the patient. Preferably, it is administered before or at the time the patient performs physical exercise.

In the present invention, it is preferred that the patient is a human. Although the present invention is applicable to both old and young people, it may find greater application in elderly people or in chronically ill people. It may also find application in people who have age-related decline in physical performance and in people who are growth hormone deficient. It is also preferred that the patient is a companion animal such as a cat or dog, with an elderly companion animal being particularly preferred.

By the term "growth hormone secretagogue" is meant any exogenously administered compound or agent that directly or indirectly stimulates or increases the endogenous release of growth hormone, growth hormone-releasing hormone or somatostatin in an animal, in particular, a human. This term shall at all times be understood to include all active forms of such secretagogues, including, for

example, the free form thereof, e.g., the free acid or base form, and also, all prodrugs, polymorphs, hydrates, solvates, stereoisomers, e.g., diastereomers and enantiomers, and the like, and all pharmaceutically acceptable salts, unless specifically stated otherwise. It will also be appreciated that suitable active
5 metabolites of growth hormone secretagogues within the scope of the present invention, in any suitable form, are also included herein.

The growth hormone secretagogue may be peptidyl or non-peptidyl in nature; however, the use of an orally active growth hormone secretagogue is preferred. In addition, it is preferred that the growth hormone secretagogue induce
10 or amplify a pulsatile release of endogenous growth hormone.


The expression "prodrug" refers to compounds that are drug precursors which, following administration, release the drug *in vivo* via some chemical or physiological process (e.g., a prodrug on being brought to the physiological pH is converted to the desired drug form). For example, a prodrug of the compound of
15 Formula I may be used in the present invention. Exemplary prodrugs are disclosed in the art, particularly in the references cited herein and incorporated herein by reference.

The growth hormone secretagogue may be used alone or in combination with another growth hormone secretagogue or with other agents which are known
20 to be useful for maintaining or increasing the benefits of exercise, such as anabolic steroids. The growth hormone secretagogue and the other agent or agents may be coadministered, either in concomitant therapy or in a fixed combination.

Representative growth hormone secretagogues are disclosed in the following International Patent Applications (listed by Publication Nos.), issued U.S.
25 patents and published European patent applications: WO 98/46569, WO 98/51687, WO 98/58947, WO 98/58949, WO 98/58950, WO 99/08697, WO 99/09991, WO 95/13069, U.S. 5,492,916, U.S. 5,494,919, WO 95/14666, WO 94/19367, WO 94/13696, WO 94/11012, U.S. 5,726,319, WO 95/11029, WO 95/17422, WO 95/17423, WO 95/34311, WO 96/02530, WO 96/22996, WO 96/22997, WO
30 96/24580, WO 96/24587, U.S. 5,559,128, WO 96/32943, WO 96/33189, WO 96/15148, WO 96/38471, WO 96/35713, WO 97/00894, WO 97/07117, WO 97/06803, WO 97/11697, WO 97/15573, WO 97/22367, WO 97/23508, WO 97/22620, WO 97/22004, WO 97/21730, WO 97/24369, U.S. 5,663,171, WO 97/34604, WO 97/36873, WO 97/40071, WO 97/40023, WO 97/41878,


WO97/41879, WO 97/46252, WO 97/44042, WO 97/38709, WO 98/03473, WO 97/43278, U.S. 5,721,251, U.S. 5,721,250, WO 98/10653, U.S. 5,919,777, U.S. 5,830,433 and EP 0995748.

- A representative first group of growth hormone secretagogues is set forth in
 5 International Patent Application, Publication No. WO 97/24369, as compounds having the structural formula below, which is designated herein as Formula II:




- wherein the various substituents are as defined in WO 97/24369. Said compounds
 10 are prepared as disclosed therein.

2-Amino-N-(2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl)-isobutyramide, having the following structure:




- 15 and 2-amino-N-(1-(R)-(2,4-difluoro-benzyloxymethyl)-2-oxo-2-(3-oxo-3a-(R)-pyridin-2-ylmethyl)-2-(2,2,2-trifluoro-ethyl)-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-ethyl)-2-methyl-propionamide, having the following structure:

-24-




and the pharmaceutically acceptable salts thereof, are within the scope of the disclosure of International Patent Application, Publication Number WO 97/24369.

A representative second group of growth hormone secretagogues is set forth in International Patent Application, Publication No. WO 98/58947, as compounds having the structural formula below, which is designated herein as Formula III:



wherein the various substituents are as defined in WO 98/58947. Said compounds are prepared as disclosed therein.

A preferred compound within this second group which may be employed in the present invention is identified as having the following name and structure: 2-amino-N-(1(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-hexahydro-imidazo[1,5-a]pyrazin-7-yl)-2-oxo-ethyl)-2-methyl-propionamide,




-25-

This compound and pharmaceutically acceptable salts thereof are within the scope of the disclosure of International Patent Application, Publication No. WO 98/58947, and may be prepared as described in Examples Five and Six therein.


A representative third group of growth hormone secretagogues is set forth in published European patent application 0995748, which discloses certain dipeptide growth hormone secretagogues of the general structural formula above, which is designated herein as Formula III, wherein the variables are as defined in EP 0995748, and their use for the treatment or prevention of musculoskeletal frailty including osteoporosis.

A representative fourth group of growth hormone secretagogues is set forth in U.S. Patent No. 5,206,235, as having the following structure:




wherein the various substituents are as defined in U.S. Patent No. 5,206,235. Said compounds are prepared as disclosed therein.

Preferred compounds within this fourth group are identified as having the following structures:




and

-26-




A representative fifth group of growth hormone secretagogues is set forth in U.S. Patent No. 5,283,241, as having the following structural formula:



- 5 wherein the various substituents are as defined in U.S. Patent 5,283,241. Said compounds are prepared as disclosed therein.

A representative sixth group of growth hormone secretagogues is disclosed in International Patent Application, Publication No. WO 97/41879, as compounds having the following structural formulas:




-27-

wherein the various substituents are as defined in WO97/41879. Said compounds are prepared as disclosed therein.

Preferred compounds within this sixth group which may be employed in the present invention are identified as having the following structure:


5



and pharmaceutically acceptable salts thereof, in particular, the methanesulfonate salt.

A representative seventh group of growth hormone secretagogues is disclosed in U.S. Patent No. 5,492,916, as being compounds of the following structural formula:

10



wherein the various substituents are as defined in U.S. Patent No. 5,492,916. Said compounds are prepared as disclosed therein.

15

All of the compounds identified above may be prepared by procedures disclosed in the cited publications. Full descriptions of the preparation of growth hormone secretagogues, which may be employed in the present invention can be found in the art, particularly in the references cited herein.


The compounds identified above and used in the present invention may have one or more asymmetric centers. The compounds of Formula I

20

72222-468

-28-

used in the present invention all have at least one asymmetric center as noted, e.g., by the asterisk in the structural Formula I-B below. Additional asymmetric centers may be present in the compounds of Formula I depending upon the nature of the various substituents on the molecule. Each such asymmetric center will produce two optical isomers and it is intended that all such optical isomers, as separated, pure or partially purified optical isomers, racemic mixtures or diastereomeric mixtures thereof, be included within the scope of the methods of the present invention. In the case of the asymmetric center represented by the asterisk, it has been found that the absolute stereochemistry of the more active and thus more preferred isomer is shown in Formula I-B below:



I-B

With the R^4 substituent as hydrogen, the spatial configuration of the asymmetric center corresponds to that in a D-amino acid. In most cases this is also designated an R-configuration although this will vary according to the values of R^3 and R^4 used in making R- or S-stereochemical assignments.

Certain compounds within the scope of the present invention may have the potential to exist in different tautomeric forms. All tautomers of a compound of the present invention are within the scope of the present invention. Also, for example, all keto-enol or imine-enamine forms of the compounds are included in the present invention. Those skilled in the art will recognize that the compound names contained herein may be based on a particular tautomer of a compound. While the name for only a particular tautomer may be used, it is intended that all tautomers are encompassed by the name of the particular tautomer and all tautomers are considered part of the present invention.

A compound within the scope of the present invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. A solvate wherein the solvent is water forms a

hydrate or hydrated ions. The present invention contemplates and encompasses both the solvated and unsolvated forms of the compounds within its scope.

Also included within the scope of the present invention are isotopically-labelled compounds, which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the present invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds of the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in compound and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of the present invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the references cited herein as well as others known in the art, by substituting a readily available isotopically labelled reagent for a non-isotopically labelled reagent.

Full descriptions of the preparation of growth hormone secretagogues employed in the present invention may be found, for example, in the following International Patent Applications (listed by Publication Nos.), issued U.S. patents and published European patent applications: WO 98/46569, WO 98/51687, WO 98/58947, WO 98/58949, WO 98/58950, WO 99/08697, WO 99/09991, WO 95/13069, U.S. 5,492,916, U.S. 5,494,919, WO 95/14666, WO 94/19367, WO 94/13696, WO 94/11012, U.S. 5,726,319, WO 95/11029, WO 95/17422, WO 95/17423, WO 95/34311, WO 96/02530, WO 96/22996, WO 96/22997, WO 96/24580, WO 96/24587, U.S. 5,559,128, WO 96/32943, WO 96/33189, WO

96/15148, WO 96/38471, WO 96/35713, WO 97/00894, WO 97/07117, WO 97/06803, WO 97/11697, WO 97/15573, WO 97/22367, WO 97/23508, WO 97/22620, WO 97/22004, WO 97/21730, WO 97/24369, U.S. 5,663,171, WO 97/34604, WO 97/36873, WO 97/40071, WO 97/40023, WO 97/41878,
5 WO97/41879, WO 97/46252, WO 97/44042, WO 97/38709, WO 98/03473, WO 97/43278, U.S. 5,721,251, U.S. 5,721,250, WO 98/10653, U.S. 5,919,777, U.S. 5,830,433 and EP 0995748.

A growth hormone secretagogue is a compound that, when administered to a patient, increases the production and/or secretion of growth hormone when
10 compared with baseline plasma concentrations of growth hormone in a normal healthy individual. Thus, to identify a growth hormone secretagogue, one need simply measure the baseline plasma concentrations of growth hormone over a time period, typically one day, and compare the plasma concentrations of growth hormone after administration of a growth hormone secretagogue with the baseline
15 concentration over the time period. Various examples of growth hormone secretagogues are disclosed herein. It is contemplated that any growth hormone secretagogue can be used in the present administration methods.

The identification of a compound as a "growth hormone secretagogue" which is able to directly or indirectly stimulate or increase the endogenous release
20 of growth hormone in an animal may be readily determined without undue experimentation by methodology well known in the art, such as the assay described by Smith et al., Science, 260, 1640-1643 (1993) (see text of Figure 2 therein). In a typical experiment, pituitary glands are aseptically removed from 150-200 g Wistar male rats and cultures of pituitary cells are prepared according to Cheng et al.,
25 Endocrinol., 124, 2791-2798 (1989). The cells are treated with the subject compound and assayed for growth hormone secreting activity, as described by Cheng et al. (ibid.). In particular, the intrinsic growth hormone secretagogue activity of a compound which may be used in the present invention may be determined by this assay.

30 The term "patient" means an animal, such as a human, a companion animal, such as a dog, cat and horse, and livestock, such as cattle, swine and sheep. Particularly preferred patients are mammals, including both males and females, with humans being even more preferred.

The term "pharmaceutically acceptable" means that a substance or mixture of substances must be compatible with the other ingredients of a formulation, and not deleterious to the patient.

5 The terms "treating", "treat" or "treatment" include preventive (e.g., prophylactic) and palliative treatment.

The term "therapeutically effective amount" means an amount of a growth hormone secretagogue that ameliorates, attenuates, or eliminates a particular disease or condition associated with growth hormone secretion and/or production, or prevents or delays the onset of a disease or condition associated with growth
10 hormone secretion and/or production.

The phrases "a compound of the present invention or a compound of Formula I" and the like, shall at all times be understood to include all active forms of such compounds, including, for example, the free form thereof, e.g., the free acid or base form, and also, all prodrugs, polymorphs, hydrates, solvates,
15 stereoisomers, e.g., diastereomers and enantiomers, and the like, and all pharmaceutically acceptable salts, unless specifically stated otherwise. It will also be appreciated that suitable active metabolites of such compounds are within the scope of the present invention.

The term "growth hormone deficient patient" (GHD patient) includes a
20 patient, such as a young adult, who has stimulated growth hormone levels below 7 ng/ml. It also includes a patient, such as an elderly patient, who does not have "clinical" growth hormone deficiency as defined above, but who is recognized as being growth hormone deficient by other measures. For example, growth hormone deficient elderly patients have average growth hormone secretion which is often as
25 much as 50 to 90% less than young adults; they have reduced average IGF-1 levels (an integrated measure of growth hormone secretion) of less than 50% of the values for young adults; they fail to have spontaneous growth hormone peaks above 5 ng/ml; and/or they have reduced sleep-associated rises in growth hormone. The decline of growth hormone secretion with aging is fully reviewed by
30 E. Corpas, S.M. Harman, M.R. Blackman, "Human Growth Hormone and Human Aging," Endocrine Review, 14:20-39 (1993).

The term "a patient with age-related decline in physical performance" is a patient who, as he or she ages, exhibits objective evidence of decline in physical performance as measured by established methods of physical performance

assessment. Measures of physical performance are objective tests of subjects' performance of standardized tasks, evaluated according to predetermined criteria that may include counting repetitions or timing the activity. A decline in physical performance results in increased risk of the patient suffering an adverse event such as an injurious fall and/or fracture. A decline in physical performance may also result in the patient having to be admitted to a nursing home and/or requiring assistance with carrying out activities of daily living.

Standardized tests of physical performance in older adults have been employed with increasing frequency in recent years. For example, J.M. Guralnik, et al., *Journal of Gerontology*, 49(2):M85-M94 (1994), and J.M. Guralnik et al., *New England Journal of Medicine*, 332:556-561 (1995), describe a short battery of physical performance tests used to assess lower extremity function in older adults. In M.E. Tinetti, et al., *Journal of the American Medical Association*, 273(17):1348-1353 (1995), physical performance skills in older adults were assessed through a series of qualitative and timed tests, described therein.

J.O. Judge et al., *Journal of the American Geriatrics Society*, 44: 1332-1341 (1996), describes measures of physical performance, such as hand grip strength, chair rise time, balance and gait speed, and found these physical performance measures were strongly associated with independence in Instrumental Activities of Daily Living (IADL). In D.B. Reuben et al., *Journal of the American Geriatrics Society*, 43: 17-23 (1995), a variety of measures of physical functioning were used, such as the Functional Status Questionnaire (FSQ), Katz Activities of Daily Living (ADL), the Older Americans Resources and Services Instrumental Activities of Daily Living (OARS-IADL), Physical Performance Test (PPT) and the Medical Outcomes Study SF-36, which were described therein. Also, workplace effectiveness tests may be used to assess one's ability to perform in the workplace.

Other physical performance tests, which are known in the art, may be used to assess physical performance. In addition, any composite scoring system, which is composed of the results from a combination of any of the above tests, may be utilized to assess physical performance.

This particular application of growth hormone secretagogues provides benefits relative to the administration of exogenous growth hormone. In particular, the growth hormone secretagogue enhances the normal pulsatile release of endogenous growth hormone and thus is more likely to reproduce the natural

pattern of endogenous growth hormone release (see J. Clin. Endocrinol. Metab. 81: 4249-4257, 1996). Growth hormone secretagogues which are orally active also have the benefit of being able to be administered orally, rather than just intravenously, intraperitoneally or subcutaneously.

5 In view of their use according to the present invention, the growth hormone secretagogues of the present invention may be formulated into various pharmaceutical forms for administration purposes. A growth hormone secretagogue may be administered, alone or in combination, by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous or subcutaneous injection, or
10 implant), nasal, vaginal, rectal, sublingual, or topical routes of administration and can be formulated with pharmaceutically acceptable carriers to provide dosage forms appropriate for each route of administration.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules and for companion animals the solid dosage forms include
15 an admixture with food and chewable forms. In such solid dosage forms, the compounds and combinations of this invention can be admixed with at least one inert pharmaceutically acceptable carrier such as sucrose, lactose, or starch. Such dosage forms can also comprise, as is normal practice, additional substances other than such inert diluents, e.g., lubricating agents such as magnesium stearate. In
20 the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings. In the case of chewable forms, the dosage form may comprise flavoring agents and perfuming agents.

Liquid dosage forms for oral administration include pharmaceutically
25 acceptable emulsions, solutions, suspensions, syrups and elixirs containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

Preparations according to this invention for parenteral administration include
30 sterile aqueous or non-aqueous solutions, suspensions, or emulsions. Examples of non-aqueous solvents or vehicles are propylene glycol, polyethylene glycol, vegetable oils, such as olive oil and corn oil, gelatin, and injectable organic esters such as ethyl oleate. Such dosage forms may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. They may be sterilized by,

for example, filtration through a bacteria-retaining filter, by incorporating sterilizing agents into the compositions, by irradiating the compositions, or by heating the compositions. They can also be manufactured in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

Compositions for rectal or vaginal administration are preferably suppositories which may contain, in addition to a compound of the present invention, excipients such as cocoa butter or a suppository wax. Compositions for nasal or sublingual administration are also prepared with standard excipients well known in the art.

The dosage of a compound of the present invention in the compositions, methods and combinations of the present invention may be varied; however, it is necessary that the amount of the compound be such that a suitable dosage form is obtained. The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment. Generally, dosage levels of between 0.0001 to 100 mg/kg of body weight daily are administered to humans and other animals, e.g., mammals, to obtain effective release of growth hormone. A preferred dosage range in humans is 0.01 to 5.0 mg/kg of body weight daily which can be administered as a single dose or divided into multiple doses.

A preferred dosage range in animals other than humans is 0.01 to 10.0 mg/kg of body weight daily, which can be administered as a single dose or divided into multiple doses. A more preferred dosage range in animals other than humans is 0.1 to 5 mg/kg of body weight daily, which can be administered as a single dose or divided into multiple doses.

Where the tartrate salt or other pharmaceutically acceptable salt of a compound of the present invention is used, the skilled person will be able to calculate effective dosage amounts by calculating the molecular weight of the salt form and performing simple stoichiometric ratios.

Also, the present invention includes within its scope the use of a growth hormone secretagogue according to the present invention, alone or in combination with another growth hormone secretagogue, such as those referenced herein, including the growth hormone releasing peptides GHRP-6 and GHRP-1 (described in U.S. Patent No. 4,411,890 and International Patent Applications, Publication

Nos. WO 89/07110, WO 89/07111), GHRP-2 (described in WO 93/04081) and B-HT920, as well as hexarelin and growth hormone releasing hormone (GHRH, also designated GRF) and its analogs, growth hormone, recombinant or natural, and its analogs and somatomedins including IGF-I and IGF-II, or in combination with other
5 therapeutic agents, such as α -adrenergic agonists such as clonidine or serotonin 5-HT1D agonists such as sumatriptan, or agents which inhibit somatostatin or its release such as physostigmine and pyridostigmine. Preferably, the growth hormone secretagogue may be used in combination with growth hormone releasing factor, an analog of growth hormone releasing factor, IGF-I or IGF-II. Also, the present
10 invention includes within its scope the use of a growth hormone secretagogue according to the present invention, alone or in combination with another therapeutic agent, such as arginine, insulin or L-dopa optionally together with propranolol.

Methods to obtain the growth hormone releasing peptides GHRP-6 and GHRP-1 are described in U.S. Patent Nos. 4,411, 890 and PCT Patent Publications
15 WO 89/07110 and WO 89/07111, methods to obtain the growth hormone releasing peptide GHRP-2 are described in PCT Patent Publication WO 93/04081, and methods to obtain hexarelin are described in J. Endocrin. Invest., 15 (Suppl. 4), 45 (1992).

In addition, the present invention includes within its scope the use of a
20 growth hormone secretagogue according to the present invention, alone or in combination with other agents, such as a growth promoting or anabolic agent, including thyrotropin-releasing hormone (TRH), parathyroid hormone (PTH), diethylstilbestrol, amino acids, estrogens, β -agonists, theophylline, anabolic steroids, enkephalins, E series prostaglandins, retinoic acid, compounds disclosed
25 in U.S. Patent No. 3,239,345, e.g., zeranol; compounds disclosed in U.S. Patent No. 4,036,979, e.g., sulbenox; or peptides disclosed in U.S. Patent No. 4,411,890.

Anabolic effects especially in the treatment of geriatric male patients are obtained with the compounds of the present invention in combination with, for example, an anabolic steroid, such as dehydroepiandrosterone (DHEA),
30 oxymetholone, methyltestosterone, fluoxymesterone, testosterone and stanozolol.

Other anabolic steroids known in the art include, for example, the following: nandrolone, primobolan, mastoron, proviron, durebolan and oxandrolone (see, for example, Histochem. Cell Biol. (2000) 113: 25-29; and J.A.M.A. 281 (14): 1282-1290 (1999)).

72222-468

-36-

For example, the active ingredient of the pharmaceutical compositions of the present invention can comprise a growth hormone secretagogue and an anabolic agent or another agent which exhibits a different activity, e.g., an antibiotic growth promoting agent, a corticosteroid to minimize catabolic side effects or
5 another pharmaceutically active material wherein the combination enhances efficacy and/or minimizes side effects, if any.

Also, the present invention includes within its scope the use of a growth hormone secretagogue according to the present invention, alone or in combination with a naturaceutic, a prodrug thereof or a pharmaceutically acceptable salt of said
10 naturaceutic or said prodrug. A naturaceutic is typically an over-the-counter composition that is promoted as, for example, improving health or general well-being. It includes compositions such as vitamins, dietary supplements, creatine, creatine phosphate and amino acids such as L-arginine.

The present invention includes within its scope the use of a pharmaceutical
15 composition according to the present invention comprising, as an active ingredient, at least one growth hormone secretagogue in association with a pharmaceutically acceptable carrier, vehicle or diluent. The present invention also includes within its scope the use of a compound of Formula I for the preparation of a medicament for the uses disclosed herein.

Combinations of these therapeutic agents, such as those mentioned herein,
20 with a growth hormone secretagogue may bring additional complementary properties to enhance the desirable properties of these various therapeutic agents. The growth hormone secretagogue and the other agent may be co-administered, either in concomitant therapy or in a fixed combination. In these combinations, the growth hormone secretagogue and the other therapeutic agent(s) may be independently present in the dose ranges from 0.01 to 1 times the dose levels
25 which are effective when these compounds and secretagogues are used singly.

Typically, the individual daily dosages for these combinations may range from about one-fifth of the minimally recommended clinical dosages to the maximum recommended levels for the entities when they are given singly. These dose ranges may be adjusted on a unit basis as necessary to permit divided daily

dosage and, as noted above, the dose will vary depending on the nature and severity of the disease, the weight of the patient, special diets and other factors.

These combinations may be formulated into pharmaceutical compositions as known in the art and as discussed herein. Since the present invention has an aspect that relates to treatment with a combination of active ingredients which may be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. The kit comprises two separate pharmaceutical compositions: a growth hormone secretagogue, a prodrug thereof or a pharmaceutically acceptable salt of said growth hormone secretagogue or said prodrug; and a second therapeutic agent as described herein. The kit comprises a container for containing the separate compositions such as a divided bottle or a divided foil packet, however, the separate compositions may also be contained within a single, undivided container. Typically, the kit comprises directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician.

An example of such a kit is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process, recesses are formed in the plastic foil. The recesses have the size and shape of the tablets or capsules to be packed. Next, the tablets or capsules are placed in the recesses and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are sealed in the recesses between the plastic foil and the sheet. Preferably, the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.

It may be desirable to provide a memory aid on the kit, e.g., in the form of numbers next to the tablets or capsules whereby the numbers correspond with the

days of the regimen which the dosage form so specified should be ingested. Another example of such a memory aid is a calendar printed on the card e.g., as follows "First Week, Monday, Tuesday, ...etc.... Second Week, Monday, Tuesday,..." etc. Other variations of memory aids will be readily apparent. A "daily dose" can be a single tablet or capsule or several tablets or capsules to be taken on a given day. Also, a daily dose of a second therapeutic agent as described herein can consist of one tablet or capsule while a daily dose of the growth hormone secretagogue, prodrug thereof or pharmaceutically acceptable salt of said growth hormone secretagogue or said prodrug can consist of several tablets or capsules or vice versa. The memory aid should reflect this.

In another specific embodiment of the invention, a dispenser designed to dispense the daily doses one at a time in the order of their intended use is provided. Preferably, the dispenser is equipped with a memory-aid, so as to further facilitate compliance with the regimen. An example of such a memory-aid is a mechanical counter which indicates the number of daily doses that has been dispensed. Another example of such a memory-aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.

The utility of the compounds described herein in the methods of the present invention are demonstrated by their activity in one or more of the assays described below.

Assay for Stimulation of Growth Hormone Release from Rat Pituitocytes

Compounds having the ability to stimulate GH secretion from cultured rat pituitary cells are identified using the following protocol. This test is also useful for comparison to standards to determine dosage levels.

Cells are isolated from pituitaries of 6-week old male Wistar rats. Following decapitation, the anterior pituitary lobes are removed into cold, sterile Hank's balanced salt solution without calcium or magnesium (HBSS). Tissues are finely minced, then subjected to two cycles of mechanically assisted enzymatic dispersion using 10 U/mL bacterial protease (EC 3.4.24.4, Sigma P-6141, St. Louis, Missouri) in HBSS. The tissue-enzyme mixture is stirred in a spinner flask at 30 rpm in a 5% CO₂ atmosphere at 37 °C for 30 min., with manual trituration after 15 min. and 30 min. using a 10-mL pipet. This mixture is centrifuged at 200 x g for 5 min. Horse

serum (35% final concentration) is added to the supernatant to neutralize excess protease. The pellet is resuspended in fresh protease (10 U/mL), stirred for about 30 min. more under the previous conditions, and manually triturated, ultimately through a 23-gauge needle. Again, horse serum (35% final concentration) is added, then the cells from both digests are combined, pelleted (200 x g for about 15 min.), resuspended in culture medium (Dulbecco's Modified Eagle Medium (D-MEM) supplemented with 4.5 g/L glucose, 10% horse serum, 2.5% fetal bovine serum, 1% non-essential amino acids, 100 U/mL nystatin and 50 mg/mL gentamycin sulfate, Gibco, Grand Island, New York) and counted. Cells are plated at $6.0\text{--}6.5 \times 10^4$ cells per cm^2 in 48-well Costar™ (Cambridge, Massachusetts) dishes and cultured for 3-4 days in culture medium.

Just prior to carrying out a GH secretion assay, culture wells are rinsed twice with release medium, then equilibrated for 30 minutes in release medium (D-MEM buffered with 25 mM Hepes, pH 7.4 and containing 0.5% bovine serum albumin at 37 °C). Test compounds are dissolved in DMSO, then diluted into pre-warmed release medium. Assays are typically run in quadruplicate. The assay is initiated by adding 0.5 mL of release medium (with vehicle or test compound) to each culture well. Incubation is carried out at 37 °C for 15 minutes, then terminated by removal of the release medium, which is centrifuged at 2000 x g for 15 minutes to remove cellular material. Rat growth hormone concentrations in the supernatants are determined by a standard radioimmunoassay protocol described below.

Assay for Exogenously-Stimulated Growth Hormone Release in the Rat after Intravenous Administration of Test Compounds

Twenty-one day old female Sprague-Dawley rats (Charles River Laboratory, Wilmington, MA) are allowed to acclimate to local vivarium conditions (24 °C, 12 hr light, 12 hr dark cycle) for approximately 1 week before testing of a compound of this invention. All rats are allowed access to water and a pelleted commercial diet (Agway Country Food, Syracuse NY) *ad libitum*.

On the day of the experiment, test compounds are dissolved in vehicle containing 1% ethanol, 1 mM acetic acid and 0.1% bovine serum albumin in saline. Each test is conducted in three rats. Rats are weighed and anesthetized via intraperitoneal injection of sodium pentobarbital (Nembutol®, 50 mg/kg body weight). Fourteen minutes after anesthetic administration, a blood sample is taken by nicking the tip of the tail and allowing the blood to drip into a microcentrifuge

-40-

tube (baseline blood sample, approximately 100 μ l). Fifteen minutes after anesthetic administration, a test compound is delivered by intravenous injection into the tail vein, with a total injection volume of 1 mL/kg body weight. Additional blood samples are taken from the tail at 5, 10 and 15 minutes after administration of a compound of this invention. Blood samples are kept on ice until serum separation by centrifugation (1430 x g for 10 minutes at 10°C). Serum is stored at -80 °C until serum growth hormone determination by radioimmunoassay as described below.

Measurement of Rat Growth Hormone

Rat growth hormone concentrations are determined by double antibody radioimmunoassay using a rat growth hormone reference preparation (NIDDK-rGH-RP-2) and rat growth hormone antiserum raised in monkey (NIDDK-anti-rGH-S-5) obtained from Dr. A. Parlow (Harbor-UCLA Medical Center, Torrance, CA). Additional rat growth hormone (1.5U/mg, #G2414, Scripps Labs, San Diego, CA) is iodinated to a specific activity of approximately 30 μ Ci/ μ g by the chloramine T method for use as tracer. Immune complexes are obtained by adding goat antiserum to monkey IgG (ICN/Cappel, Aurora, OH) plus polyethylene glycol, MW 10,000-20,000 to a final concentration of 4.3%; recovery is accomplished by centrifugation according to methods well known to those skilled in the art. This assay has a working range of 0.08-2.5 μ g rat growth hormone per tube.

Assessment of Growth Hormone Release in the Dog after Oral Administration

On the day of dosing, the test compound is weighed out for the appropriate dose and dissolved in water. Doses are delivered at a volume of 0.5-3 mL/kg by oral gavage to 2-4 dogs for each dosing regimen. Blood samples (5 mL) are collected from the jugular vein by direct venipuncture pre-dose and at 0.17, 0.33, 0.5, 0.75, 1, 2, 4, 6, 8 and 24 hours post dose using 5 mL vacutainers containing lithium heparin. The prepared plasma is stored at -20 °C until analysis.

Measurement of Canine Growth Hormone

Canine growth hormone concentrations are determined by a standard radioimmunoassay protocol using canine growth hormone (antigen for iodination and reference preparation AFP-1983B) and canine growth hormone antiserum raised in monkey (AFP-21452578) obtained from Dr. A. Parlow (Harbor-UCLA Medical Center, Torrance, CA). Tracer is produced by chloramine T-iodination of canine growth hormone to a specific activity of 20-40 μ Ci/ μ g. Immune complexes

are obtained by adding goat antiserum to monkey IgG (ICN/Cappel, Aurora, OH) plus polyethylene glycol, MW 10,000-20,000 to a final concentration of 4.3%; recovery is accomplished by centrifugation according to methods well known to those skilled in the art. This assay has a working range of 0.08-2.5 µg canine
5 GH/tube.

Phase I Double-Blind, Placebo-Controlled, Multiple Dose Evaluation
of Test Compound in Calorically Restricted, Obese Men and Women

A total of 48 obese subjects, without significant other medical problems and a negative exercise tolerance test, were recruited for this study and placed on a
10 caloric-restricted diet. Subjects had been randomized to either placebo, 3 mg po tid, 6 mg po tid or 10 mg po qam of the test compound, 2-amino-N-[2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl]-isobutyramide, L-tartrate for a total of seven days.

One of the objectives of this study was to examine the effect of the test
15 compound on exercise-induced growth hormone secretion. The testing was conducted on an exercise bicycle. The results of this objective of this study are summarized in the table below.

<u>Treatment</u>	<u>N</u>	<u>Before Treatment</u>		<u>End of Treatment</u>		<u>Change</u>		<u>p-value</u> ¹
		<u>Mean</u>	<u>SD</u>	<u>Mean</u>	<u>SD</u>	<u>Mean</u>	<u>SD</u>	
Placebo	10	126	134	194	168	68	56	
3 mg p.o. tid	10	87	80	384	295	298	329	0.0146
6 mg p.o. tid	12	145	149	395	203	250	176	0.0056
10 mg p.o. qam	10	243	235	927	374	684	315	<0.0001

1. 2-sample Wilcoxon test comparing the active change with the placebo change

2. SD represents standard deviation

One of the efficacy measures of this study was the area under the growth hormone secretory response curve over time (AUC). The above table summarizes the growth hormone AUC values at the screening visit ("Before Treatment") and at Day 7 ("End of Treatment") for subjects, who completed exercise at both visits. Also, the AUC values for these subjects were calculated for up to 150 minutes post exercise. The "p-value" in the above table, which is associated with each active treatment group, is derived from comparing each active treatment group with the placebo group, using the two-sample Wilcoxon test. In this study, it was found that the test compound augmented exercise-induced growth hormone secretion.

While the foregoing description discloses the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the present invention encompasses all of the usual variations, adaptations or modifications as come within the scope of the following claims and their equivalents.

72222-468

-43-

CLAIMS:

1. A drug for increasing physical exercise-stimulated growth hormone secretion in a patient, which comprises:

(a) a therapeutically effective amount of a growth hormone secretagogue, and

(b) a pharmaceutically acceptable carrier.

2. A drug for increasing improvement in muscle mass or strength produced by physical exercise in a patient or for increasing improvement in physical performance produced by physical exercise in a patient, which comprises:

(a) a therapeutically effective amount of a growth hormone secretagogue, and

(b) a pharmaceutically acceptable carrier,

wherein the drug is adapted to be taken in conjunction with physical exercise by the patient.

3. A drug for increasing physical exercise capacity in a patient in need thereof, which comprises:

(a) a therapeutically effective amount of a growth hormone secretagogue, and

(b) a pharmaceutically acceptable carrier,

wherein the drug is adapted to be taken in conjunction with physical exercise by the patient.

72222-468

-43a-

4. A drug for maintaining or improving protein balance in a patient, which comprises:

(a) a therapeutically effective amount of a growth hormone secretagogue, and

5 (b) a pharmaceutically acceptable carrier,

wherein the drug is adapted to be taken with physical exercise by the patient.

5. A drug for preservation of improvement in muscle mass or strength in a patient or for preservation of
10 improvement in physical performance in a patient, which comprises:

(a) a therapeutically effective amount of a growth hormone secretagogue, and

(b) a pharmaceutically acceptable carrier,

15 wherein the drug is adapted to be taken between periods of physical exercise by the patient.

6. A drug for treating sarcopenia in a patient in need thereof, which comprises:

(a) a therapeutically effective amount of a growth
20 hormone secretagogue, and

(b) a pharmaceutically acceptable carrier,

wherein the drug is adapted to be taken in conjunction with physical exercise by the patient.


7. The drug of claim 1, 2, 3, 4, 5 or 6, wherein the
25 physical exercise is physical rehabilitation, physical therapy or is part of a regular physical exercise program.

72222-468

-43b-

8. The drug of claim 1, 2, 3, 4, 5, 6 or 7, wherein the patient is a human who is elderly, chronically ill, obese, has age-related decline in physical performance or is growth hormone deficient.
- 5 9. The drug of any one of claims 1 to 8, wherein the growth hormone secretagogue is a compound of Formula I:





-44-




I

- or a stereoisomeric mixture thereof, diastereomerically enriched, diastereomerically pure, enantiomerically enriched or enantiomerically pure isomer thereof, or a
 5 prodrug of such compound, mixture or isomer thereof, or a pharmaceutically acceptable salt of the compound, mixture, isomer or prodrug, or a tautomer thereof, wherein:

HET is a heterocyclic moiety selected from the group consisting of



and



- 10 d is 0, 1 or 2;
 e is 1 or 2;
 f is 0 or 1;
 n and w are 0, 1 or 2, provided that n and w cannot both be 0 at the same time;
 Y² is oxygen or sulfur;

A is a divalent radical, where the left hand side of the radical as shown below is connected to C" and the right hand side of the radical as shown below is connected to C', selected from the group consisting of

- 5 -NR²-C(O)-NR²-, -NR²-S(O)₂-NR²-, -O-C(O)-NR²-, -NR²-C(O)-O-, -C(O)-NR²-C(O)-, -C(O)-NR²-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-NR²-C(O)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-,
 -S(O)₂-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-O-C(O)-, -C(R⁹R¹⁰)-O-C(R⁹R¹⁰)-,
 -NR²-C(O)-C(R⁹R¹⁰)-, -O-C(O)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(O)-NR²-,
 -C(R⁹R¹⁰)-C(O)-O-, -C(O)-NR²-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(O)-O-C(R⁹R¹⁰)-,
 -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -S(O)₂-NR²-C(R⁹R¹⁰)-C(R⁹R¹⁰)-,
 10 -C(R⁹R¹⁰)-C(R⁹R¹⁰)-NR²-C(O)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-O-C(O)-,
 -NR²-C(O)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -NR²-S(O)₂-C(R⁹R¹⁰)-C(R⁹R¹⁰)-,
 -O-C(O)-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(O)-NR²-,
 -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(O)-, -C(R⁹R¹⁰)-NR²-C(O)-O-, -C(R⁹R¹⁰)-O-C(O)-NR²-,
 -C(R⁹R¹⁰)-NR²-C(O)-NR²-, -NR²-C(O)-O-C(R⁹R¹⁰)-, -NR²-C(O)-NR²-C(R⁹R¹⁰)-,
 15 -NR²-S(O)₂-NR²-C(R⁹R¹⁰)-, -O-C(O)-NR²-C(R⁹R¹⁰)-, -C(O)-N=C(R¹¹)-NR²-,
 -C(O)-NR²-C(R¹¹)=N-, -C(R⁹R¹⁰)-NR¹²-C(R⁹R¹⁰)-, -NR¹²-C(R⁹R¹⁰)-,
 -NR¹²-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(O)-O-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -NR²-C(R¹¹)=N-C(O)-,
 -C(R⁹R¹⁰)-C(R⁹R¹⁰)-N(R¹²)-, -C(R⁹R¹⁰)-NR¹²-, -N=C(R¹¹)-NR²-C(O)-,
 -C(R⁹R¹⁰)-C(R⁹R¹⁰)-NR²-S(O)₂-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-S(O)₂-NR²-,
 20 -C(R⁹R¹⁰)-C(R⁹R¹⁰)-C(O)-O-, -C(R⁹R¹⁰)-S(O)₂-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-S(O)₂-,
 -O-C(R⁹R¹⁰)-C(R⁹R¹⁰)-, -C(R⁹R¹⁰)-C(R⁹R¹⁰)-O-, -C(R⁹R¹⁰)-C(O)-C(R⁹R¹⁰)-,
 -C(O)-C(R⁹R¹⁰)-C(R⁹R¹⁰)- and -C(R⁹R¹⁰)-NR²-S(O)₂-NR²-;

Q is a covalent bond or CH₂;

W is CH or N;

- 25 X is CR⁹R¹⁰, C=CH₂ or C=O;

Y is CR⁹R¹⁰, O or NR²;

Z is C=O, C=S or S(O)₂;

- G¹ is hydrogen, halo, hydroxy, nitro, amino, cyano, phenyl, carboxyl, -CONH₂, -(C₁-C₄)alkyl optionally independently substituted with one or more phenyl, one or more
 30 halogens or one or more hydroxy groups, -(C₁-C₄)alkoxy optionally independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, -(C₁-C₄)alkylthio, phenoxy, -COO(C₁-C₄)alkyl, N,N-di-(C₁-C₄)alkylamino, -(C₂-C₆)alkenyl optionally independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, -(C₂-C₆)alkynyl optionally

independently substituted with one or more phenyl, one or more halogens or one or more hydroxy groups, $-(C_3-C_6)$ cycloalkyl optionally independently substituted with one or more (C_1-C_4) alkyl groups, one or more halogens or one or more hydroxy groups, $-(C_1-C_4)$ alkylamino carbonyl or di- (C_1-C_4) alkylamino carbonyl;

- 5 G^2 and G^3 are each independently selected from the group consisting of hydrogen, halo, hydroxy, $-(C_1-C_4)$ alkyl optionally independently substituted with one to three halo groups and $-(C_1-C_4)$ alkoxy optionally independently substituted with one to three halo groups;

- R^1 is hydrogen, $-CN$, $-(CH_2)_qN(X^6)C(O)X^6$, $-(CH_2)_qN(X^6)C(O)(CH_2)_t-A^1$,
 10 $-(CH_2)_qN(X^6)S(O)_2(CH_2)_t-A^1$, $-(CH_2)_qN(X^6)S(O)_2X^6$, $-(CH_2)_qN(X^6)C(O)N(X^6)(CH_2)_t-A^1$,
 $-(CH_2)_qN(X^6)C(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)N(X^6)(X^6)$, $-(CH_2)_qC(O)N(X^6)(CH_2)_t-A^1$,
 $-(CH_2)_qC(O)OX^6$, $-(CH_2)_qC(O)O(CH_2)_t-A^1$, $-(CH_2)_qOX^6$, $-(CH_2)_qOC(O)X^6$,
 $-(CH_2)_qOC(O)(CH_2)_t-A^1$, $-(CH_2)_qOC(O)N(X^6)(CH_2)_t-A^1$, $-(CH_2)_qOC(O)N(X^6)(X^6)$,
 $-(CH_2)_qC(O)X^6$, $-(CH_2)_qC(O)(CH_2)_t-A^1$, $-(CH_2)_qN(X^6)C(O)OX^6$,
 15 $-(CH_2)_qN(X^6)S(O)_2N(X^6)(X^6)$, $-(CH_2)_qS(O)_mX^6$, $-(CH_2)_qS(O)_m(CH_2)_t-A^1$,
 $-(C_1-C_{10})$ alkyl, $-(CH_2)_t-A^1$, $-(CH_2)_q-(C_3-C_7)$ cycloalkyl, $-(CH_2)_q-Y^1-(C_1-C_6)$ alkyl,
 $-(CH_2)_q-Y^1-(CH_2)_t-A^1$ or $-(CH_2)_q-Y^1-(CH_2)_t-(C_3-C_7)$ cycloalkyl;

where the alkyl and cycloalkyl groups in the definition of R^1 are optionally substituted with (C_1-C_4) alkyl, hydroxy, (C_1-C_4) alkoxy, carboxyl, $-CONH_2$,

- 20 $-S(O)_m(C_1-C_6)$ alkyl, $-CO_2(C_1-C_4)$ alkyl ester, 1H-tetrazol-5-yl or 1, 2 or 3 fluoro groups;

Y^1 is O, $S(O)_m$, $-C(O)NX^6-$, $-CH=CH-$, $-C\equiv C-$, $-N(X^6)C(O)-$, $-C(O)NX^6-$,
 $-C(O)O-$, $-OC(O)N(X^6)-$ or $-OC(O)-$;

q is 0, 1, 2, 3 or 4;

- 25 t is 0, 1, 2 or 3;

said $(CH_2)_q$ group and $(CH_2)_t$ group in the definition of R^1 are optionally independently substituted with hydroxy, (C_1-C_4) alkoxy, carboxyl, $-CONH_2$,

$-S(O)_m(C_1-C_6)$ alkyl, $-CO_2(C_1-C_4)$ alkyl ester, 1H-tetrazol-5-yl, 1, 2 or 3 fluoro groups or 1 or 2 (C_1-C_4) alkyl groups;

- 30 R^{1A} is selected from the group consisting of hydrogen, F, Cl, Br, I, (C_1-C_6) alkyl, phenyl (C_1-C_3) alkyl, pyridyl (C_1-C_3) alkyl, thiazolyl (C_1-C_3) alkyl and thienyl (C_1-C_3) alkyl, provided that R^{1A} is not F, Cl, Br or I when a heteroatom is vicinal to C^* ;

R^2 is hydrogen, (C_1-C_8) alkyl, $-(C_0-C_3)$ alkyl- (C_3-C_8) cycloalkyl, $-(C_1-C_4)$ alkyl- A^1 or A^1 ;

-47-

where the alkyl groups and the cycloalkyl groups in the definition of R^2 are optionally substituted with hydroxy, $-C(O)OX^6$, $-C(O)N(X^6)(X^6)$, $-N(X^6)(X^6)$, $-S(O)_m(C_1-C_6)alkyl$, $-C(O)A^1$, $-C(O)(X^6)$, CF_3 , CN or 1, 2 or 3 independently selected halo groups;

- 5 R^3 is selected from the group consisting of A^1 , $(C_1-C_{10})alkyl$, $-(C_1-C_6)alkyl-A^1$, $-(C_1-C_6)alkyl-(C_3-C_7)cycloalkyl$, $-(C_1-C_5)alkyl-X^1-(C_1-C_5)alkyl$, $-(C_1-C_5)alkyl-X^1-(C_0-C_5)alkyl-A^1$ and $-(C_1-C_5)alkyl-X^1-(C_1-C_5)alkyl-(C_3-C_7)cycloalkyl$;

where the alkyl groups in the definition of R^3 are optionally substituted with $-S(O)_m(C_1-C_6)alkyl$, $-C(O)OX^3$, 1, 2, 3, 4 or 5 independently selected halo groups or 1, 2 or 3 independently selected $-OX^3$ groups;


10

X^1 is O, $S(O)_m$, $-N(X^2)C(O)-$, $-C(O)N(X^2)-$, $-OC(O)-$, $-C(O)O-$, $-CX^2=CX^2-$, $-N(X^2)C(O)O-$, $-OC(O)N(X^2)-$ or $-C\equiv C-$;

- R^4 is hydrogen, $(C_1-C_6)alkyl$ or $(C_3-C_7)cycloalkyl$, or R^4 is taken together with R^3 and the carbon atom to which they are attached and form $(C_5-C_7)cycloalkyl$, $(C_5-C_7)cycloalkenyl$, a partially saturated or fully saturated 4- to 8-membered ring having 1 to 4 heteroatoms independently selected from the group consisting of oxygen, sulfur and nitrogen, or is a bicyclic ring system consisting of a partially saturated or fully saturated 5- or 6-membered ring, fused to a partially saturated, fully unsaturated or fully saturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

20

X^4 is hydrogen or $(C_1-C_6)alkyl$ or X^4 is taken together with R^4 and the nitrogen atom to which X^4 is attached and the carbon atom to which R^4 is attached and form a five to seven membered ring;



where a and b are each independently 0, 1, 2 or 3;

X^5 and X^{5a} are each independently selected from the group consisting of hydrogen, CF_3 , A^1 and optionally substituted $(C_1-C_6)alkyl$;

30

the optionally substituted $(C_1-C_6)alkyl$ in the definition of X^5 and X^{5a} is optionally substituted with a substituent selected from the group consisting of A^1 , OX^2 , $-S(O)_m(C_1-C_6)alkyl$, $-C(O)OX^2$, $(C_3-C_7)cycloalkyl$, $-N(X^2)(X^2)$ and $-C(O)N(X^2)(X^2)$;

-48-

or the carbon bearing X^5 or X^{5a} forms one or two alkylene bridges with the nitrogen atom bearing R^7 and R^8 wherein each alkylene bridge contains 1 to 5 carbon atoms, provided that when one alkylene bridge is formed then only one of X^5 or X^{5a} is on the carbon atom and only one of R^7 or R^8 is on the nitrogen atom and further provided that when two alkylene bridges are formed then X^5 and X^{5a} cannot be on the carbon atom and R^7 and R^8 cannot be on the nitrogen atom;


or X^5 is taken together with X^{5a} and the carbon atom to which they are attached and form a partially saturated or fully saturated 3- to 7-membered ring, or a partially saturated or fully saturated 4- to 8-membered ring having 1 to 4 heteroatoms independently selected from the group consisting of oxygen, sulfur and nitrogen;

or X^5 is taken together with X^{5a} and the carbon atom to which they are attached and form a bicyclic ring system consisting of a partially saturated or fully saturated 5- or 6-membered ring, optionally having 1 or 2 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated, fully saturated or fully unsaturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

Z^1 is a bond, O or $N-X^2$, provided that when a and b are both 0 then Z^1 is not $N-X^2$ or O;

or R^6 is $-(CR^aR^b)_a-E-(CR^aR^b)_b-$, where the $-(CR^aR^b)_a-$ group is attached to the carbonyl carbon of the amide group of the compound of formula I and the $-(CR^aR^b)_b$ group is attached to the terminal nitrogen atom of the compound of formula I;

E is -O-, -S-, -CH=CH- or an aromatic moiety selected from



said aromatic moiety in the definition of E optionally substituted with up to three halo, hydroxy, $-N(R^c)(R^c)$, (C_1-C_6) alkyl or (C_1-C_6) alkoxy;

R^a and R^b are, for each occurrence, independently hydrogen, (C_1-C_6) alkyl, trifluoromethyl, phenyl or monosubstituted (C_1-C_6) alkyl where the

5 substituents are imidazolyl, naphthyl, phenyl, indolyl, p-hydroxyphenyl, $-OR^c$, $S(O)_mR^c$, $C(O)OR^c$, (C_3-C_7) cycloalkyl, $-N(R^c)(R^c)$, $-C(O)N(R^c)(R^c)$, or R^a or R^b may independently be joined to one or both of R^7 or E (where E is other than O, S or $-CH=CH-$) to form an alkylene bridge between the terminal nitrogen and the alkyl portion of the R^a or R^b and the R^7 or E group,

10 wherein the bridge contains 1 to 8 carbon atoms; or R^a and R^b may be joined to one another to form a (C_3-C_7) cycloalkyl;

R^c , for each occurrence, is independently hydrogen or (C_1-C_6) alkyl;

a and b are independently 0, 1, 2 or 3, with the proviso that if E is $-O-$ or $-S-$, b is other than 0 or 1 and with the further proviso that if E is $-CH=CH-$, b

15 is other than 0;

R^7 and R^8 are each independently hydrogen or optionally substituted (C_1-C_6) alkyl; where the optionally substituted (C_1-C_6) alkyl in the definition of R^7 and R^8 is optionally independently substituted with A^1 , $-C(O)O-(C_1-C_6)$ alkyl, $-S(O)_m(C_1-C_6)$ alkyl, 1 to 5 halo groups, 1 to 3 hydroxy groups, 1 to 3

20 $-O-C(O)(C_1-C_{10})$ alkyl groups or 1 to 3 (C_1-C_6) alkoxy groups; or R^7 and R^8 can be taken together to form $-(CH_2)_r-L-(CH_2)_r-$; where L is $C(X^2)(X^2)$, $S(O)_m$ or $N(X^2)$;

R^9 and R^{10} are each independently selected from the group consisting of hydrogen, fluoro, hydroxy and (C_1-C_5) alkyl optionally independently substituted with 1-5 halo

25 groups;

R^{11} is selected from the group consisting of (C_1-C_5) alkyl and phenyl optionally substituted with 1-3 substituents each independently selected from the group consisting of (C_1-C_5) alkyl, halo and (C_1-C_5) alkoxy;

R^{12} is selected from the group consisting of (C_1-C_5) alkylsulfonyl, (C_1-C_5) alkanoyl

30 and (C_1-C_5) alkyl where the alkyl portion is optionally independently substituted by 1-5 halo groups;

A^1 for each occurrence is independently selected from the group consisting of (C_5-C_7) cycloalkenyl, phenyl, a partially saturated, fully saturated or fully unsaturated 4- to 8-membered ring optionally having 1 to 4 heteroatoms independently selected

from the group consisting of oxygen, sulfur and nitrogen and a bicyclic ring system consisting of a partially saturated, fully unsaturated or fully saturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated, fully saturated or fully unsaturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

A¹ for each occurrence is independently optionally substituted, on one or optionally both rings if A¹ is a bicyclic ring system, with up to three substituents, each substituent independently selected from the group consisting of F, Cl, Br, I, OCF₃, OCF₂H, CF₃, CH₃, OCH₃, -OX⁶, -C(O)N(X⁶)(X⁶), -C(O)OX⁶, oxo, (C₁-C₆)alkyl, nitro, cyano, benzyl, -S(O)_m(C₁-C₆)alkyl, 1H-tetrazol-5-yl, phenyl, phenoxy, phenylalkyloxy, halophenyl, methylenedioxy, -N(X⁶)(X⁶), -N(X⁶)C(O)(X⁶), -S(O)₂N(X⁶)(X⁶), -N(X⁶)S(O)₂-phenyl, -N(X⁶)S(O)₂X⁶, -CONX¹¹X¹², -S(O)₂NX¹¹X¹², -NX⁶S(O)₂X¹², -NX⁶CONX¹¹X¹², -NX⁶S(O)₂NX¹¹X¹², -NX⁶C(O)X¹², imidazolyl, thiazolyl and tetrazolyl, provided that if A¹ is optionally substituted with methylenedioxy then it can only be substituted with one methylenedioxy;

where X¹¹ is hydrogen or optionally substituted (C₁-C₆)alkyl; the optionally substituted (C₁-C₆)alkyl defined for X¹¹ is optionally independently substituted with phenyl, phenoxy, (C₁-C₆)alkoxycarbonyl, -S(O)_m(C₁-C₆)alkyl, 1 to 5 halo groups, 1 to 3 hydroxy groups, 1 to 3 (C₁-C₁₀)alkanoyloxy groups or 1 to 3 (C₁-C₆)alkoxy groups;

X¹² is hydrogen, (C₁-C₆)alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl, provided that when X¹² is not hydrogen, the X¹² group is optionally substituted with one to three substituents independently selected from the group consisting of Cl, F, CH₃, OCH₃, OCF₃ and CF₃;

or X¹¹ and X¹² are taken together to form -(CH₂)_r-L¹-(CH₂)_r;

L¹ is C(X²)(X²), O, S(O)_m or N(X²);

r for each occurrence is independently 1, 2 or 3;

X² for each occurrence is independently hydrogen, optionally substituted (C₁-C₆)alkyl or optionally substituted (C₃-C₇)cycloalkyl, where the optionally substituted

72222-468

-51-

(C₁-C₆)alkyl and optionally substituted (C₃-C₇)cycloalkyl in the definition of X² are optionally independently substituted with -S(O)_m(C₁-C₆)alkyl, -C(O)OX³, 1 to 5 halo groups or 1-3 OX³ groups;

X³ for each occurrence is independently hydrogen or (C₁-C₆)alkyl;

- 5 X⁶ for each occurrence is independently hydrogen, optionally substituted (C₁-C₆)alkyl, (C₂-C₆)halogenated alkyl, optionally substituted (C₃-C₇)cycloalkyl, (C₃-C₇)halogenated cycloalkyl, where optionally substituted (C₁-C₆)alkyl and optionally substituted (C₃-C₇)cycloalkyl in the definition of X⁶ is optionally independently mono- or di-substituted with (C₁-C₄)alkyl, hydroxy, (C₁-C₄)alkoxy, carboxyl, CONH₂,
 10 -S(O)_m(C₁-C₆)alkyl, carboxylate (C₁-C₄)alkyl ester or 1H-tetrazol-5-yl; or
 when there are two X⁶ groups on one atom and both X⁶ are independently (C₁-C₆)alkyl, the two (C₁-C₆)alkyl groups may be optionally joined and, together with the atom to which the two X⁶ groups are attached, form a 4- to 9- membered ring optionally having oxygen, sulfur or NX⁷ as a ring member;


- 15 X⁷ is hydrogen or (C₁-C₆)alkyl optionally substituted with hydroxy;

m for each occurrence is independently 0, 1 or 2;

with the provisos that:

- 1) X⁶ and X¹² cannot be hydrogen when attached to C(O) or S(O)₂ in the form C(O)X⁶, C(O)X¹², S(O)₂X⁶ or S(O)₂X¹²; and
 20 2) when R⁶ is a bond then L is N(X²) and each r in the definition -(CH₂)_r-L-(CH₂)_r- is independently 2 or 3.

10. The drug of claim 9 wherein the growth hormone secretagogue is a compound of Formula I-A.




- 25 a racemic-diastereomeric mixture or an optical isomer of the compound or a pharmaceutically-acceptable salt or a prodrug thereof, or a tautomer thereof,
 wherein
 f is 0;
 n is 0 and w is 2, or n is 1 and w is 1, or n is 2 and w is 0;
 30 Y is oxygen or sulfur;

- R^1 is hydrogen, $-\text{CN}$, $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{C}(\text{O})\text{X}^6$, $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{C}(\text{O})(\text{CH}_2)_t\text{A}^1$,
 $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{SO}_2(\text{CH}_2)_t\text{A}^1$, $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{SO}_2\text{X}^6$, $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{C}(\text{O})\text{N}(\text{X}^6)(\text{CH}_2)_t\text{A}^1$,
 $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{C}(\text{O})\text{N}(\text{X}^6)(\text{X}^6)$, $-(\text{CH}_2)_q\text{C}(\text{O})\text{N}(\text{X}^6)(\text{X}^6)$, $-(\text{CH}_2)_q\text{C}(\text{O})\text{N}(\text{X}^6)(\text{CH}_2)_t\text{A}^1$,
 $-(\text{CH}_2)_q\text{C}(\text{O})\text{OX}^6$, $-(\text{CH}_2)_q\text{C}(\text{O})\text{O}(\text{CH}_2)_t\text{A}^1$, $-(\text{CH}_2)_q\text{OX}^6$, $-(\text{CH}_2)_q\text{OC}(\text{O})\text{X}^6$,
5 $-(\text{CH}_2)_q\text{OC}(\text{O})(\text{CH}_2)_t\text{A}^1$, $-(\text{CH}_2)_q\text{OC}(\text{O})\text{N}(\text{X}^6)(\text{CH}_2)_t\text{A}^1$, $-(\text{CH}_2)_q\text{OC}(\text{O})\text{N}(\text{X}^6)(\text{X}^6)$,
 $-(\text{CH}_2)_q\text{C}(\text{O})\text{X}^6$, $-(\text{CH}_2)_q\text{C}(\text{O})(\text{CH}_2)_t\text{A}^1$, $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{C}(\text{O})\text{OX}^6$,
 $-(\text{CH}_2)_q\text{N}(\text{X}^6)\text{SO}_2\text{N}(\text{X}^6)(\text{X}^6)$, $-(\text{CH}_2)_q\text{S}(\text{O})_m\text{X}^6$, $-(\text{CH}_2)_q\text{S}(\text{O})_m(\text{CH}_2)_t\text{A}^1$,
 $-(\text{C}_1\text{-C}_{10})\text{alkyl}$, $-(\text{CH}_2)_t\text{A}^1$, $-(\text{CH}_2)_q\text{-(C}_3\text{-C}_7\text{)cycloalkyl}$, $-(\text{CH}_2)_q\text{-Y}^1\text{-(C}_1\text{-C}_6\text{)alkyl}$,
 $-(\text{CH}_2)_q\text{-Y}^1\text{-(CH}_2)_t\text{A}^1$ or $-(\text{CH}_2)_q\text{-Y}^1\text{-(CH}_2)_t\text{-(C}_3\text{-C}_7\text{)cycloalkyl}$;
10 where the alkyl and cycloalkyl groups in the definition of R^1 are optionally
substituted with $(\text{C}_1\text{-C}_4)\text{alkyl}$, hydroxyl, $(\text{C}_1\text{-C}_4)\text{alkoxy}$, carboxyl, $-\text{CONH}_2$,
 $-\text{S}(\text{O})_m(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{CO}_2(\text{C}_1\text{-C}_4)\text{alkyl ester}$, 1H-tetrazol-5-yl or 1, 2 or 3
fluoro;
 Y^1 is O, $\text{S}(\text{O})_m$, $-\text{C}(\text{O})\text{NX}^6$ -, $-\text{CH}=\text{CH}$ -, $-\text{C}\equiv\text{C}$ -, $-\text{N}(\text{X}^6)\text{C}(\text{O})$ -, $-\text{C}(\text{O})\text{O}$ -,
15 $-\text{OC}(\text{O})\text{N}(\text{X}^6)$ - or $-\text{OC}(\text{O})$ -;
 q is 0, 1, 2, 3 or 4;
 t is 0, 1, 2 or 3;
said $(\text{CH}_2)_q$ group and $(\text{CH}_2)_t$ group may each be optionally substituted with
hydroxyl, $(\text{C}_1\text{-C}_4)\text{alkoxy}$, carboxyl, $-\text{CONH}_2$, $-\text{S}(\text{O})_m(\text{C}_1\text{-C}_6)\text{alkyl}$,
20 $-\text{CO}_2(\text{C}_1\text{-C}_4)\text{alkyl ester}$, 1H-tetrazol-5-yl, 1, 2 or 3 fluoro, or 1 or 2 $(\text{C}_1\text{-C}_4)\text{alkyl}$;
 R^2 is hydrogen, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-(\text{C}_0\text{-C}_3)\text{alkyl-(C}_3\text{-C}_6\text{)cycloalkyl}$, $-(\text{C}_1\text{-C}_4)\text{alkyl-A}^1$ or A^1 ;
where the alkyl groups and the cycloalkyl groups in the definition of R^2 are
optionally substituted with hydroxyl, $-\text{C}(\text{O})\text{OX}^6$, $-\text{C}(\text{O})\text{N}(\text{X}^6)(\text{X}^6)$,
25 $-\text{N}(\text{X}^6)(\text{X}^6)$, $-\text{S}(\text{O})_m(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C}(\text{O})\text{A}^1$, $-\text{C}(\text{O})(\text{X}^6)$, CF_3 , CN or 1, 2 or 3
halogen;
 R^3 is A^1 , $(\text{C}_1\text{-C}_{10})\text{alkyl}$, $-(\text{C}_1\text{-C}_6)\text{alkyl-A}^1$, $-(\text{C}_1\text{-C}_6)\text{alkyl-(C}_3\text{-C}_7\text{)cycloalkyl}$,
 $-(\text{C}_1\text{-C}_5)\text{alkyl-X}^1\text{-(C}_1\text{-C}_5)\text{alkyl}$, $-(\text{C}_1\text{-C}_5)\text{alkyl-X}^1\text{-(C}_0\text{-C}_5)\text{alkyl-A}^1$ or
 $-(\text{C}_1\text{-C}_5)\text{alkyl-X}^1\text{-(C}_1\text{-C}_5)\text{alkyl-(C}_3\text{-C}_7\text{)cycloalkyl}$;
30 where the alkyl groups in the definition of R^3 are optionally substituted with,
 $-\text{S}(\text{O})_m(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C}(\text{O})\text{OX}^3$, 1, 2, 3, 4 or 5 halogens, or 1, 2 or 3 OX^3 ;
 X^1 is O, $\text{S}(\text{O})_m$, $-\text{N}(\text{X}^2)\text{C}(\text{O})$ -, $-\text{C}(\text{O})\text{N}(\text{X}^2)$ -, $-\text{OC}(\text{O})$ -, $-\text{C}(\text{O})\text{O}$ -, $-\text{CX}^2=\text{CX}^2$ -,
 $-\text{N}(\text{X}^2)\text{C}(\text{O})\text{O}$ -, $-\text{OC}(\text{O})\text{N}(\text{X}^2)$ - or $-\text{C}\equiv\text{C}$;
 R^4 is hydrogen, $(\text{C}_1\text{-C}_6)\text{alkyl}$ or $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$;

-53-

X^4 is hydrogen or (C_1-C_6) alkyl or X^4 is taken together with R^4 and the nitrogen atom to which X^4 is attached and the carbon atom to which R^4 is attached and form a five to seven membered ring;



5 where a and b are independently 0, 1, 2 or 3;

X^5 and X^{5a} are each independently selected from the group consisting of hydrogen, trifluoromethyl, A^1 and optionally substituted (C_1-C_6) alkyl;

the optionally substituted (C_1-C_6) alkyl in the definition of X^5 and X^{5a} is optionally substituted with a substituent selected from the group consisting of A^1 , OX^2 , $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^2$,
10 (C_3-C_7) cycloalkyl, $-N(X^2)(X^2)$ and $-C(O)N(X^2)(X^2)$;

R^7 and R^8 are independently hydrogen or optionally substituted (C_1-C_6) alkyl;

where the optionally substituted (C_1-C_6) alkyl in the definition of R^7 and R^8 is optionally independently substituted with A^1 , $-C(O)O-(C_1-C_6)$ alkyl,

15 $-S(O)_m(C_1-C_6)$ alkyl, 1 to 5 halogens, 1 to 3 hydroxy, 1 to 3 $-O-C(O)(C_1-C_{10})$ alkyl or 1 to 3 (C_1-C_6) alkoxy; or

R^7 and R^8 can be taken together to form $-(CH_2)_r-L-(CH_2)_r-$;

where L is $C(X^2)(X^2)$, $S(O)_m$ or $N(X^2)$;

A^1 in the definition of R^1 is a partially saturated, fully saturated or fully unsaturated
20 4- to 8-membered ring optionally having 1 to 4 heteroatoms independently selected from the group consisting of oxygen, sulfur and nitrogen, a bicyclic ring system consisting of a partially saturated, fully unsaturated or fully saturated 5- or 6-membered ring, having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated, fully
25 saturated or fully unsaturated 5- or 6-membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

A^1 in the definition of R^2 , R^3 , R^6 , R^7 and R^8 is independently (C_5-C_7) cycloalkenyl, phenyl or a partially saturated, fully saturated or fully unsaturated 4- to 8-
30 membered ring optionally having 1 to 4 heteroatoms independently selected from the group consisting of oxygen, sulfur and nitrogen, a bicyclic ring system consisting of a partially saturated, fully unsaturated or fully saturated 5- or 6-

membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen, fused to a partially saturated, fully saturated or fully unsaturated 5- or 6- membered ring, optionally having 1 to 4 heteroatoms independently selected from the group consisting of nitrogen, sulfur and oxygen;

A^1 for each occurrence is independently optionally substituted, in one or optionally both rings if A^1 is a bicyclic ring system, with up to three substituents, each substituent independently selected from the group consisting of F, Cl, Br, I, OCF_3 , OCF_2H , CF_3 , CH_3 , OCH_3 , $-OX^6$, $-C(O)N(X^6)(X^6)$, $-C(O)OX^6$, oxo, (C_1-C_6) alkyl, nitro, cyano, benzyl, $-S(O)_m(C_1-C_6)$ alkyl, 1H-tetrazol-5-yl, phenyl, phenoxy, phenylalkyloxy, halophenyl, methylenedioxy, $-N(X^6)(X^6)$, $-N(X^6)C(O)(X^6)$, $-SO_2N(X^6)(X^6)$, $-N(X^6)SO_2$ -phenyl, $-N(X^6)SO_2X^6$, $-CONX^{11}X^{12}$, $-SO_2NX^{11}X^{12}$, $-NX^6SO_2X^{12}$, $-NX^6CONX^{11}X^{12}$, $-NX^6SO_2NX^{11}X^{12}$, $-NX^6C(O)X^{12}$, imidazolyl, thiazolyl or tetrazolyl, provided that if A^1 is optionally substituted with methylenedioxy then it can only be substituted with one methylenedioxy;

where X^{11} is hydrogen or optionally substituted (C_1-C_6) alkyl;

the optionally substituted (C_1-C_6) alkyl defined for X^{11} is optionally independently substituted with phenyl, phenoxy, (C_1-C_6) alkoxycarbonyl, $-S(O)_m(C_1-C_6)$ alkyl 1 to 5 halogens, 1 to 3 hydroxy, 1 to 3 (C_1-C_{10}) alkanoyloxy or 1 to 3 (C_1-C_6) alkoxy;

X^{12} is hydrogen, (C_1-C_6) alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl, provided that when X^{12} is not hydrogen, X^{12} is optionally substituted with one to three substituents independently selected from the group consisting of Cl, F, CH_3 , OCH_3 , OCF_3 and CF_3 ;

or X^{11} and X^{12} are taken together to form $-(CH_2)_rL^1-(CH_2)_r$;

where L^1 is $C(X^2)(X^2)$, O, $S(O)_m$ or $N(X^2)$;

r for each occurrence is independently 1, 2 or 3;

X^2 for each occurrence is independently hydrogen, optionally substituted (C_1-C_6) alkyl, or optionally substituted (C_3-C_7) cycloalkyl, where the optionally substituted (C_1-C_6) alkyl and optionally substituted (C_3-C_7) cycloalkyl in the definition of X^2 are optionally independently substituted with $-S(O)_m(C_1-C_6)$ alkyl, $-C(O)OX^3$, 1 to 5 halogens or 1-3 OX^3 ;

72222-468

-55-

X^3 for each occurrence is independently hydrogen or (C₁-C₆)alkyl;

X^6 is independently hydrogen, optionally substituted (C₁-C₆)alkyl, (C₂-C₆)halogenated alkyl, optionally substituted (C₃-C₇)cycloalkyl, (C₃-C₇)halogenatedcycloalkyl, where optionally substituted (C₁-C₆)alkyl and optionally substituted (C₃-C₇)cycloalkyl in the definition of X^6 is optionally independently substituted by 1 or 2 (C₁-C₄)alkyl, hydroxyl, (C₁-C₄)alkoxy, carboxyl, CONH₂, -S(O)_m(C₁-C₆)alkyl, carboxylate (C₁-C₄)alkyl ester, or 1H-tetrazol-5-yl; or when there are two X^6 groups on one atom and both X^6 are independently (C₁-C₆)alkyl, the two (C₁-C₆)alkyl groups may be optionally joined and, together with the atom to which the two X^6 groups are attached, form a 4- to 9- membered ring optionally having oxygen, sulfur or NX⁷;

X^7 is hydrogen or (C₁-C₆)alkyl optionally substituted with hydroxyl; and

m for each occurrence is independently 0, 1 or 2;

with the proviso that:

X^6 and X^{12} cannot be hydrogen when it is attached to C(O) or SO₂ in the form C(O) X^6 , C(O) X^{12} , SO₂ X^6 or SO₂ X^{12} ; and when R^6 is a bond then L is N(X^2) and each r in the definition -(CH₂)_r-L-(CH₂)_r- is independently 2 or 3.

11. The drug of claim 10 wherein the compound, the prodrug thereof or the pharmaceutically acceptable salt of the compound or the prodrug, is selected from the group consisting of:

2-amino-N-(2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl)-isobutyramide;

2-amino-N-[2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl]-isobutyramide, L-tartrate;

2-amino-N-(1-(R)-(2,4-difluoro-benzyloxymethyl)-2-oxo-2-(3-oxo-3a-(R)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-ethyl)-2-methyl-propionamide; and

the (L)-(+)-tartaric acid salt of 2-amino-N-(1-(R)-(2,4-difluoro-benzyloxymethyl)-2-oxo-2-(3-oxo-3a-(R)-pyridin-2-ylmethyl-2-(2,2,2-trifluoro-ethyl)-2,3,3a,4,6,7-hexahydro-pyrazolo-[4,3-c]pyridin-5-yl)-ethyl)-2-methyl-propionamide.

72222-468

-56-

12. The drug of claim 1 wherein the growth hormone secretagogue is selected from the group consisting of:

2-amino-N-{1(R)-benzyloxymethyl-2-[1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl]-2-oxoethyl}-2-methylpropionamide; and

the (L)-(+)-tartaric acid salt of 2-amino-N-(1(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethyl)-2-methylpropionamide.

13. The drug of claim 2 wherein the growth hormone secretagogue is selected from the group consisting of:

2-amino-N-{1(R)-benzyloxymethyl-2-[1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl]-2-oxoethyl}-2-methylpropionamide; and

the (L)-(+)-tartaric acid salt of 2-amino-N-(1(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethyl)-2-methylpropionamide.

14. The drug of claim 3 wherein the growth hormone secretagogue is selected from the group consisting of:

2-amino-N-{1(R)-benzyloxymethyl-2-[1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl]-2-oxoethyl}-2-methylpropionamide; and

the (L)-(+)-tartaric acid salt of 2-amino-N-(1(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-

72222-468

-57-

(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethyl)-2-methylpropionamide.

15. The drug of claim 4 wherein the growth hormone secretagogue is selected from the group consisting of:

5 2-amino-N-{1(R)-benzyloxymethyl-2-[1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl]-2-oxoethyl}-2-methylpropionamide; and

the (L)-(+)-tartaric acid salt of 2-amino-N-(1(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethyl)-2-methylpropionamide.

16. The drug of claim 5 wherein the growth hormone secretagogue is selected from the group consisting of:

15 2-amino-N-{1(R)-benzyloxymethyl-2-[1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl]-2-oxoethyl}-2-methylpropionamide; and

the (L)-(+)-tartaric acid salt of 2-amino-N-(1(R)-benzyloxymethyl-2-(1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl)-2-oxoethyl)-2-methylpropionamide.

17. The drug of claim 6 wherein the growth hormone secretagogue is selected from the group consisting of:

25 2-amino-N-{1(R)-benzyloxymethyl-2-[1,3-dioxo-8a(S)-pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl]-2-oxoethyl}-2-methylpropionamide; and

72222-468

-58-

the (L)-(+) -tartaric acid salt of 2-amino-N-(1(R) - benzyloxymethyl-2-(1,3-dioxo-8a(S) -pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl) - 2-oxoethyl) -2-methylpropionamide.

- 5 18. The drug of claim 7 wherein the growth hormone secretagogue is selected from the group consisting of:

2-amino-N-{1(R) -benzyloxymethyl-2-[1,3-dioxo-8a(S) -pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl] -2-
10 oxoethyl} -2-methylpropionamide; and

the (L)-(+) -tartaric acid salt of 2-amino-N-(1(R) - benzyloxymethyl-2-(1,3-dioxo-8a(S) -pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl) - 2-oxoethyl) -2-methylpropionamide.

- 15 19. The drug of claim 8 wherein the growth hormone secretagogue is selected from the group consisting of:

2-amino-N-{1(R) -benzyloxymethyl-2-[1,3-dioxo-8a(S) -pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl] -2-
20 oxoethyl} -2-methylpropionamide; and

the (L)-(+) -tartaric acid salt of 2-amino-N-(1(R) - benzyloxymethyl-2-(1,3-dioxo-8a(S) -pyridin-2-ylmethyl-2-(2,2,2-trifluoroethyl)hexahydroimidazo[1,5-a]pyrazin-7-yl) - 2-oxoethyl) -2-methylpropionamide.

- 25 20. The drug of any one of claims 1 to 19, which further comprises:

(c) a recombinant growth hormone or a growth hormone secretagogue selected from the group consisting of GHRP-6, GHRP-1, GHRP-2, hexarelin, growth hormone releasing

72222-468

-59-

factor, an analog of growth hormone releasing factor, IGF-I and IGF-II.

21. The drug of any one of claims 1 to 19, which further comprises:

5 (d) an anabolic steroid, a prodrug thereof or a pharmaceutically acceptable salt of the steroid or the prodrug.

22. The drug of any one of claims 1 to 19, which further comprises:

10 (e) a naturaceutic, a prodrug thereof or a pharmaceutically acceptable salt of the naturaceutic or the prodrug.

23. The drug of any one of claims 12 to 19, which is adapted to be taken by a human and contains the ingredient

15 (a) at a daily dosage of 0.01 to 5.0 mg/kg body weight.

24. A commercial package comprising the drug as defined in claim 12 and a written matter which is associated therewith and describes indications of the drug for use in a human for increasing physical exercise-stimulated growth
20 hormone secretion.

25. A commercial package comprising the drug as defined in claim 13 and a written matter which is associated therewith and describes indications of the drug for use in a human for increasing improvement in muscle mass or strength
25 produced, by physical exercise by taking the drug in conjunction with the physical exercise.

26. A commercial package comprising the drug as defined in claim 13 and a written matter which is associated therewith and describes indications of the drug for use in a

72222-468

-60-

human for increasing improvement in physical performance produced by physical exercise, by taking the drug in conjunction with the physical exercise.

27. A commercial package comprising the drug as
5 defined in claim 14 and a written matter which is associated therewith and describes indications of the drug for use in a human for increasing physical exercise capacity, by taking the drug in conjunction with the physical exercise.

28. A commercial package comprising the drug as
10 defined in claim 15 and a written matter which is associated therewith and describes indications of the drug for use in a human for maintaining or improving protein balance with physical exercise, by taking the drug in conjunction with the physical exercise.

15 29. A commercial package comprising the drug as defined in claim 16 and a written matter which is associated therewith and describes indications of the drug for use in a human for preservation of improvement in muscle mass or strength by physical exercise, by taking the drug in
20 conjunction with the physical exercise.

30. A commercial package comprising the drug as defined in claim 16 and a written matter which is associated therewith and describes indications of the drug for use in a human for preservation of improvement in physical
25 performance, by taking the drug in conjunction with the physical exercise.

31. A commercial package comprising the drug as defined in claim 17 and a written matter which is associated therewith and describes indications of the drug for use in a
30 human for treating sarcopenia, by taking the drug with physical exercise.

72222-468

-61-

32. The commercial package of any one of claims 24 to 31, wherein the human is obese.

33. The commercial package of claim 32, wherein the physical exercise is part of a regular physical exercise
5 program.

34. The commercial package of any one of claims 24 to 31, wherein the human is elderly, chronically ill or growth hormone deficient, or has age-related decline in physical performance.

10 35. The commercial package of claim 34, wherein the physical exercise is physical rehabilitation or physical therapy.

SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS